



Effects of High Intensity Interval Training on Plasma Levels of GH and IGF-I Seyyed Mahmoud Hejazi*

Department of Physical Education, Mashhad Branch, Islamic Azad University, Mashhad, Iran

*Corresponding e-mail: sm.hejazi37@gmail.com

ABSTRACT

Introduction: It is well-recognized that exercise has a significant impact on the GH/IGF system but less is known about the effects of HIIT on this axis. Aim of the present study was to evaluate the effect of ten weeks of HIIT on plasma levels of GH and IGF-I in healthy men. **Methods:** Twenty young men (age 23.34 ± 2.56 weight 72.47 ± 12.01 height 174.10 ± 5.75) recruited and randomly assigned into control ($n=10$) and HIIT ($n=12$) groups. HIIT protocol was started with 4 cycles. Then, every two weeks one cycle was added to the previous ones. Finally, it was to 8 cycles/session in tenth weeks that lasted 16 minutes. Blood samples were collected prior to and after HIIT program for all subjects and IGF-I and GH levels were measured. **Result:** HIIT subjects showed a significant increase in IGF-I ($P=0.002$, $F=12.38$). However, no significant change was shown in GH levels ($P=0.716$, $F=0.62$). **Discussion and conclusion:** Our findings indicate that the HIIT caused increase in circulating levels of IGF-I independently from GH levels. Both hormones may contribute to positive effects of anabolic conditions.

Keywords: IGF-I/GH axis, HIIT, adaptation

INTRODUCTION

Growth hormone (GH) is the principal regulator of the hepatic synthesis of insulin-like growth factor I (IGF-1). IGF-1 itself is the primary downstream mediator of GH actions, and circulating IGF-1 plays an important role in the feedback regulation of GH secretion [1]. However, IGF-1, produced in skeletal muscle during exercise, is also released into the circulation which might explain an increase in Circulating IGF-1 levels as well [2].

IGF-1 has widespread anabolic and insulin-sensitizing effects, and plays a critical role in formation, maintenance, and regeneration of skeletal muscles [1]. IGF-1 also plays a direct role in whole body glucose homeostasis primarily by stimulating skeletal muscle glucose uptake [3]. Nindl, et al. stated that IGF-1 is an important metabolic biomarker associated with a variety of health and exercise-related outcomes [2].

It is well-recognized that exercise has a significant impact on the GH/IGF system. number of factors have led to interest in the effect of exercise on the growth hormone/insulin-like growth factor-I (GH/IGF-I) axis, including its possible role in maintenance of lean mass [4]. In a variety of physiological actions (protein synthesis, cellular proliferation, glucose metabolism, etc.) [5].

Although most modes of exercise stimulate an increased GH secretory response that is linear with exercise intensity [6] evidence suggests IGF-I responses are independent of GH [7].

Insulin-like growth factor I (IGF-1) is a polypeptide of 70 amino acids (7650 Daltons), and is one of a number of related insulin-like growth factors present in the circulation. The molecule has a number of biological activities similar to insulin. IGF-1 concentrations change with age, nutritional status, body composition and physical activity [8].

Whether previous studies have reported exercise-induced alterations of IGF-I seems to depend on several factors, including exercise model. Both low- and high-intensity cycling have been shown to increase IGF-I concentrations [9].

However, neither low-volume nor high-volume resistance exercise has been shown to change total IGF-I concentrations [10].

Moreover, no change in IGF-I concentrations has been found following a marathon, a 20 km run, and treadmill exercise at 60% of $V_{o\max}$ [11,12].

The ability of IGF-I to promote muscle hypertrophy is unchallenged; however, several lines of evidence have demonstrated that load-induced hypertrophy can occur independently of IGF-I and/or activation of the IGF-I receptor [13]. Conversely, evidence in support of IGF-I as a “regulator or amplifier” of muscle remodelling cascades also exists [14].

HIIT exercises are high intensity and interval that can be done in a short time, although has the benefits of long-term endurance exercise [15]. Even though the recent studies have shown endurance training can induce an increase of GH, IGF-1, levels in the circulation, but less is known about the effects of different training intensities (e. g. high-intensity training (HIIT) on circulating levels of these growth factors [3]. Only a few studies have addressed this issue in a strictly experimental way [16]. The purpose of the present study was to evaluate the effect of ten weeks of HIIT on plasma levels of GH and IGF-I in healthy men. The present study differs from previous studies, the protocol was more rigorous and exercise responses were compared with a non-exercise control trial.

METHODS

Study design/participants

Trial design was semi-experimental with control group. Twenty-two young men (age 23.34 ± 2.56 weight 72.47 ± 12.01 height 174.10 ± 5.75) recruited via a recall in Ferdowsi university of Mashhad campuses and those approved participations were randomized into either a training group (HIIT) or a control group (CON). Informed consent was obtained from each patient included in the study and the study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki.

Exclusion criteria include professional athletics history as well as the current regular exercise, smoking, cardiovascular and metabolic disease, or any complication that disrupt the implementation of exercise.

High intensity interval training protocol

HIIT group after became acquainted with the correct training performances, carried out the exercises, every other day, three sessions a week, for 10 weeks. Exercises included warm up, HIIT training, cool down. Subjects warmed up by stretching and easy walking for 5 minutes.

The HIIT interventions consisted of four 30 s maximal effort (“all-out”) shuttle run bouts (from cone-1 to cone-2, 20-meter sweep) separated by 1.5 min passive rest each (Figure 1).




Figure 1 High intensity interval training protocol

HIIT protocol began with 4 cycles and every two weeks added one cycle. Finally, it was 8 cycles in tenth week that lasted 16 minutes. Cooling Down also included 5 min of walking and stretching. During the HIIT interventions, all subjects were vocally encouraged to maintain Maximal effort. Quality of training was controlled by a physical education expert and subject’s heartbeat was constantly checked by polar device. Control group asked to be sedentary in this period.

Blood sampling/Hormonal measurement

Twenty-four hours before starting the exercise program, while all the subjects were fastened; 5 cc of blood was taken from their brachial vein. Also, sampling repeated after 48 hours of last session in same condition. Blood samples were

frozen at -20°C for future analysis. LDN (Germany) ELISA assay kit was used to measure IGF-I concentrations which sensitivity was 1.292 ng/mL and Monobind (USA) ELISA assay kit was used to measure GH concentrations which Sensitivity was 0.072 ng/mL.

Statistical Analysis

Data normality was tested using Shapiro-wilk and the homogeneity of the variances was tested using levene. After making sure of the normality and equality of the groups, variance analysis with repeated measuring was used to study the differences between groups. Statistical analysis was done by the SPSS Software.

RESULTS

After ten weeks HIIT subjects showed a significant increase in IGF-I ($P=0.002$, $F=12.38$). However, no significant change was shown in GH levels ($P=0.716$, $F=0.62$) (Table 1).

Table 1: Study variables before (pre) and after (post) 10 weeks of study protocol

Variable	Control		HIIT		ANOVA (Repeated Measurement)
	Pre-test	Post-test	Pre-test	Post-test	
IGF-I (ng /mL)	193.10 \pm 33.54	190.60 \pm 32.32	199.27 \pm 28.69	222.27 \pm 32.90	F=12.38 P=0.002
GH (ng /mL)	3.053 \pm 0.73	3.004 \pm 0.53	3.241 \pm 0.41	3.158 \pm 0.61	F=0.62 P=0.716

Data are mean \pm SD

DISCUSSION AND CONCLUSION

Previous studies have been reported that exercise has a significant impact on the levels of several hormones, and can increase resistance and performance, as well as muscle mass. Hormone levels can change according to several parameters, including the type and length of exercise, the duration of time following exercise, the age and gender of the athletes, among others [17].

Based on the findings from the present research, plasma levels of IGF-I had been significantly increased due to HIIT protocol. To our knowledge, no previous research has directly investigated the impacts of HIIT on IGF-I/GH axis. However, other types of training have been published. Most of the training time during the HIIT intervention was spent in recovery between short, intense bursts of all-out shuttle running. This is in accordance with previously published studies [18].

Several studies have indicated that anabolic hormones, such as insulin, GH, testosterone and IGF-1, stimulate neural tissue and muscle development during resistance exercise [19].

The serum concentration of anabolic hormones is elevated during and following resistance exercise compared to the level at rest, which leads to hypertrophy and remodelling of muscles [20]. Circadian rhythm has specific effects on the release of IGF-1 in the body, where the hormone levels are higher in the morning and lower in the afternoon [21].

In the present study, GH concentrations did not significantly change as a result of relative long term HIIT. That regard should be considered that due to our study limitation, GH only measured at one point in time. Also, regarding Circadian rhythm and the pulsatile manner of GH, it will probably cause different results when compared with multi-time point.

whereas long-term exercise training approximately doubles integrated GH concentrations when measured on non-exercising days [22]. Linnamo, et al. reports that GH levels are increased in response to submaximal and maximal heavy resistance exercise. However, the prominent increase was detected just after the exercise session was completed, and the response returned to normal level two hours post exercise [23].

Different training intensities, such as high-intensity training and high volume, low-intensity training may have a different impact on hormone levels. Although pH is generally well regulated, a more increase in the acidity of the circulating blood and the skeletal muscle occurs when performing HIIT. One can speculate that these systemic and local changes in the extracellular environment might influence the release, the affinity, and association/dissociation of GH, IGF-1. The extracellular pH has been recognized to regulate the IGF-1 interactions with different cells, components of the extracellular [24,25].

In summary, we undertook a randomized trial of the impact of 10 weeks of HIIT on IGF-I/GH axis in previously untrained subjects. The major finding was HIIT caused increase in circulating levels of IGF-I independently from GH levels. Both hormones may contribute to the positive effects of anabolic conditions as it has been shown by previous studies [26].

REFERENCES

- [1] Frystyk, Jan. "Exercise and the growth hormone-insulin-like growth factor axis." *Medicine and science in sports and exercise* 42.1 (2010): 58-66.
- [2] Nindl, Bradley C. "Insulin-like growth factor-I, physical activity, and control of cellular anabolism." *Medicine & Science in Sports & Exercise* 42.1 (2010): 35-38.
- [3] Berg, Ulrika, and Peter Bang. "Exercise and circulating insulin-like growth factor I." *Hormone Research in Paediatrics* 62.S1 (2005): 50-58.
- [4] Brill, Kimberly T., et al. "Single and combined effects of growth hormone and testosterone administration on measures of body composition, physical performance, mood, sexual function, bone turnover, and muscle gene expression in healthy older men." *The Journal of Clinical Endocrinology & Metabolism* 87.12 (2002): 5649-5657.
- [5] Weltman, Arthur, et al. "Impact of abdominal visceral fat, growth hormone, fitness, and insulin on lipids and lipoproteins in older adults." *Metabolism* 52.1 (2003): 73-80.
- [6] Pritzlaff-Roy, Cathy J., et al. "Gender governs the relationship between exercise intensity and growth hormone release in young adults." *Journal of Applied Physiology* 92.5 (2002): 2053-2060.
- [7] Hornum, Mette, et al. "Exercise-induced changes in circulating growth factors with cyclic variation in plasma estradiol in women." *Journal of Applied Physiology* 82.6 (1997): 1946-1951.
- [8] Stitt, Trevor N., et al. "The IGF-1/PI3K/Akt pathway prevents expression of muscle atrophy-induced ubiquitin ligases by inhibiting FOXO transcription factors." *Molecular cell* 14.3 (2004): 395-403.
- [9] Cappon, James, et al. "Effect of brief exercise on circulating insulin-like growth factor I." *Journal of Applied Physiology* 76.6 (1994): 2490-2496.
- [10] Nindl, Bradley C., et al. "Overnight responses of the circulating IGF-I system after acute, heavy-resistance exercise." *Journal of Applied Physiology* 90.4 (2001): 1319-1326.
- [11] Hagberg, James M., et al. "Metabolic responses to exercise in young and older athletes and sedentary men." *Journal of Applied Physiology* 65.2 (1988): 900-908.
- [12] Banfi, G., et al. "Growth hormone and insulin-like growth factor I in athletes performing a marathon at 4000 m of altitude." *Growth regulation* 4.2 (1994): 82-86.
- [13] Spangenburg, Espen E., et al. "A functional insulin-like growth factor receptor is not necessary for load-induced skeletal muscle hypertrophy." *The Journal of physiology* 586.1 (2008): 283-291.
- [14] Flueck, Martin, and Geoffrey Goldspink. "Counterpoint: IGF is not the major physiological regulator of muscle mass." *Journal of Applied Physiology* 108.6 (2010): 1821-1823.
- [15] Laursen, Paul B., and David G. Jenkins. "The scientific basis for high-intensity interval training." *Sports Medicine* 32.1 (2002): 53-73.
- [16] Schwarz, Adam J., et al. "Acute effect of brief low-and high-intensity exercise on circulating insulin-like growth factor (IGF) I, II, and IGF-binding protein-3 and its proteolysis in young healthy men." *The Journal of Clinical Endocrinology & Metabolism* 81.10 (1996): 3492-3497.
- [17] Kraemer, Robert R., et al. "Similar hormonal responses to concentric and eccentric muscle actions using relative loading." *European journal of applied physiology* 96.5 (2006): 551-557.
- [18] Wahl, Patrick, et al. "Effect of high-and low-intensity exercise and metabolic acidosis on levels of GH, IGF-I, IGFBP-3 and cortisol." *Growth Hormone & IGF Research* 20.5 (2010): 380-385.
- [19] Crewther, Blair, et al. "Possible stimuli for strength and power adaptation." *Sports medicine* 36.3 (2006): 215-238.
- [20] Widdowson, W. Matthew, et al. "The physiology of growth hormone and sport." *Growth Hormone & IGF Research* 19.4 (2009): 308-319.
- [21] Hayes, Lawrence D., Gordon F. Bickerstaff, and Julien S. Baker. "Interactions of cortisol, testosterone, and resistance training: influence of circadian rhythms." *Chronobiology international* 27.4 (2010): 675-705.

- [22] Weltman, Arthur, et al. "Endurance training amplifies the pulsatile release of growth hormone: Effects of training intensity." *Journal of Applied Physiology* 72.6 (1992): 2188-2196.
- [23] Linnamo, Vesa, et al. "Acute hormonal responses to submaximal and maximal heavy resistance and explosive exercises in men and women." *The Journal of Strength & Conditioning Research* 19.3 (2005): 566-571.
- [24] Forsten-Williams, Kimberly, et al. "Enhanced insulin-like growth factor-I (IGF-I) cell association at reduced pH is dependent on IGF binding protein-3 (IGFBP-3) interaction." *Journal of cellular physiology* 210.2 (2007): 298-308.
- [25] Gordon, Scott E., et al. "Effect of acid-base balance on the growth hormone response to acute high-intensity cycle exercise." *Journal of Applied Physiology* 76.2 (1994): 821-829.
- [26] Gibala, Martin J., et al. "Short-term sprint interval versus traditional endurance training: Similar initial adaptations in human skeletal muscle and exercise performance." *The Journal of physiology* 575.3 (2006): 901-911.