



International Journal of Medical Research & Health Sciences

www.ijmrhs.com

Volume 4 Issue 3

Coden: IJMRRS

Copyright ©2015

ISSN: 2319-5886

Received: 17th Mar 2015Revised: 20th Apr 2015Accepted: 29th May 2015

Research article

REGENERATION OF ARTICULAR CARTILAGE UNDER THE IMPLANTATION OF BONE MATRIX

Yuri M. Iryanov¹, *Nikolay A. Kiryanov², Olga V. Dyuriagina³, Tatiana Yu. Karaseva⁴, Evgenii A. Karasev⁵

¹Head of the Laboratory of Morphology, Doctor of Biological Sciences, Professor. Russian Ilizarov Scientific Center Restorative Traumatology and Orthopaedics (RISC RTO), Kurgan, Russia

²Head of Department of Pathology of Izhevsk State Medical Academy, Doctor of Medical Science, Professor, Russia

³A Senior Researcher, Laboratory of Purulent Osteology and Limb Defect Filling, RISC RTO, Candidate of Veterinary Sciences, Russia

⁴Head, ⁵A Researcher, Department of Orthopaedics, RISC RTO, Candidate of Medical Sciences, Russia

*Corresponding author email: kirknik@list.ru

ABSTRACT

Background: The damage or loss of articular cartilage is costly medical problem. The purpose of this work - morphological analysis of reparative chondrogenesis when implanted in the area of the knee joint cartilage of granulated mineralized bone matrix. **Material and Methods:** The characteristic features of the knee cartilage regeneration studied experimentally in pubertal Wistar rats after modeling a marginal perforated defect and implantation of granulated mineralized bone matrix obtained according to original technology without heat and demineralizing processing into the injury zone. **Results:** This biomaterial established to have pronounced chondro- and osteoinductive properties, and to provide prolonged activation of reparative process, accelerated organotypical remodeling and restoration of the articular cartilage injured. **Conclusion:** The data obtained demonstrate the efficacy of in clinical practice for the treatment of diseases and injuries of the articular cartilage.

Keywords: Articular cartilage, The mineral component of bone matrix, Implantation, Reparative chondrogenesis.

INTRODUCTION

The damage or loss of the articular cartilage due to congenital anomalies, degenerative joint diseases or injuries has a negative effect on the quality of life of the injured in all age-related groups, and it is expensive to treat ^[1]. Surgical methods are often inadequate, and in many cases they do not give positive results in terms of proper replacement of articular hyaline cartilage ^[2, 3]. In this regard, the development, scientific substantiation, and experimental morphologic evaluation of the biomaterials which have chondrogenic activity are of major importance in modern medicine. The most widely used biomaterials (demineralized bone matrix, matrices of polylactic and polyglycolic acids /PLA

and PGA/, collagen cryogels, analogs of bone mineral, polysaccharides of natural origin) are characterized by low plastic effectiveness, the absence of chondrogenic activity, and limited biocompatibility ^[4, 5, 6,7]. Our previous studies have demonstrated that mineralized bone matrix (MBM) obtained by using original technology without thermal and demineralizing processing ^[8] has osteoinductive properties, and it stimulates osteogenesis effectively ^[9]. The research is aimed at performing morphological analysis of reparative chondrogenesis when this biomaterial is implanted into the zone of the knee cartilage injury.

MATERIAL AND METHODS

All experiments in this study were performed in accordance with guidelines for animal research and were approved by our academy Ethics Committee. Experimental groups. The experiments were conducted on 25 pubertal Wistar rats male of 340–390 g weight (20 experimental and 5 intact ones). The process of keeping, experimenting, and euthanizing animals was regulated by the normative documents [2]. The operations on the animals were performed in an operating room under general anesthesia (8 mg of Rometar and 4 mg of Zoletil per 1000 g of body weight intramuscularly). Non-through fenestrated defects of 2.5-3 mm in diameter were made on the patellar surface of the distal left and right femur with the dental drill, the subchondral bone was penetrated. Sterile granulated allogenic MBM (2-3-mg) obtained from tubular bones of mongrel rats by original technology [6] was placed into the right bone defect. The defect in the left bone was left untreated under blood clot and it was used as control. The knee cartilage was studied in five intact normal rats for comparison. The animals were taken out of the experiment after 7, 15, 30, and 60 days after the surgery (five animals were used for each time point).

Morphological evaluation. The pieces of femur bones of the operated rats and those of intact ones were fixed and embedded in paraffin (after decalcification) and in araldite (without decalcification). Paraffin sections were stained with hematoxylin-eosin, with picrofuchsin according to Van Gieson, and with Alcian blue (Alcian blue 8GS) at pH 2.5 and 1.0 to reveal non-sulphated glycosaminoglycans (NSGAG) and sulphated ones (SGAG), respectively. The content of glycosaminoglycans was determined in conventional units (c.u.) by defining Alcian blue concentration in sections which was measured using INCA-200 Energy energy-dispersion X-ray spectrometer (X-ray electron probe microanalyzer) (Oxford instruments, England) [4]. The structure of MBM granules and regenerated bones in the injured area was investigated using JSM-840 scanning electron microscope and JEM-2010 transmission electron microscope (Jeol, Japan). The results of quantitative investigations were processed using the methods of variation statistics. The significance of the differences of the parameters compared was

calculated using Student's t-test. The differences were considered significant at the level of <0.05 .

RESULTS

The implanted MBM granules are of 50-200 μm in diameter and have a well-ordered highly porous structure. The pores correspond to the places of localizing bone lacunae and bone tubules from which cells and other organic components have been removed when obtaining the biomaterial.

Seven days after the surgery, the joint capsule in both left and right limbs was hyperemic and edematic. The cartilaginous coating was dull. The signs of inflammatory reaction, and alternatively destructive changes affecting all metaepiphyseal components could be revealed around the injury. Foci of hematoma were found. They were infiltrated by fibrin clots, little-differentiated cell elements, neutrophilic granulocytes, macrophages, mast cells, extravasal erythrocytes, and lymphocytes. Leukocytic necrotic masses containing lysed cells and fibrin strata could be observed. Articular cartilage was scarified. The defect zone was partially replenished by granular and little-differentiated loose edematic connective tissue and few vessels. Sporadic bone-osteoid foci and isolated thin bone trabeculae were identified around the injury of the right bone (experiment). Reparative bone formation occurred by the type of intramembranous osteogenesis. NSGAG and SGAG content in the zone of articular cartilage injury was reduced significantly comparing with the similar values in intact animals (0.59 ± 0.03 , and 0.68 ± 0.03 c.u., respectively), however, there were no significant changes revealed in their content in control and experiment at this stage (Table 1).

Table 1: Glycosaminoglycane content in the zone of the knee cartilage injury in rats without defect plasty (control), and when using allogenic MBM (experiment) ($M\pm m$, c.u.)

The period of the experiment, days	Control		Experiment	
	NSGAG	SGAG	NSGAG	SGAG
7	0.18 ± 0.01	0.11 ± 0.01	0.20 ± 0.01	0.12 ± 0.01
15	0.19 ± 0.01	0.12 ± 0.01	$0.49\pm 0.01^*$	$0.26\pm 0.01^*$
30	0.21 ± 0.01	0.16 ± 0.01	$0.56\pm 0.02^*$	$0.54\pm 0.01^*$
60	0.22 ± 0.01	0.24 ± 0.01	$0.59\pm 0.03^*$	$0.70\pm 0.01^*$

*Significant changes comparing with control values. Student's t-test <0.05

Fifteen days after the surgery, numerous leukocytes, fibroblastic elements, bundles of collagen fibers, granulation tissue, and a significant number of dilated and filled with blood vessels could be identified in the defect central zone of control bone. Cellular-and-fibrillar elements of inflammation were not found in the right bone (experiment), reparative bone formation occurred not only by the type of intramembranous osteogenesis, but by that of enchondral osteogenesis as well. In the zone of subchondral bone injury an extensive small-looped network was formed consisting of thickened trabeculae covered by rows of numerous large osteoblasts which produced spongy bone closely adhered with the surface of underlying bone. Implanted MBM granules resembled cystic cavities formed during decalcification of the sample and due to biodestruction. Functionally active osteoclasts with multiple nuclei and a brush-like border were located within these cavities (Fig. 1). Numerous sinusoids were surrounded by proliferating peri-vasculocytes.




Fig 1: Zone of the defect of the rat knee cartilage 15 days after the surgery: a – n osteoclast within the implanted MBM granule which has the form of a cystic cavity, paraffin sections, staining with hematoxylin-eosin, magnification 400; b –

ultrastructure of the osteoclast localized on the surface of the implanted MBM granule, the arrow points to the osteoclast brush-like border in the zone of contact, ultrathin sections, transmission electron microscopy, magnification 8000.

There were no hemorrhages and destruction foci in these areas, as well as active proliferation of fibroblasts, intense neoangiogenesis was observed. The layers of osteo- and chondrogenic cells at different differentiation stages were located on the granule surface. The zone of articular cartilage defect was partially filled with newly formed hyaline cartilage. Its intercellular substance contained NSGAG predominantly (Table 1). The defect edges merged with the edges of “maternal” articular cartilage where cell nests with isogenic groups were identified. The presence of the latter was evidenced by proliferative activity of some chondrocytes which were one of the sources of the cartilaginous tissue filling the defect. The cells located round MBM granules were the second source, and the chondrocytes in the zone of enchondral osteogenesis in the sites of subchondral bone fracture healing presented the third one. Glycosaminoglycane concentration in the defect zone for experiment more than twice exceeded control values (Table 1).

A significant part of bone defect was filled with loose or dense connective tissue 30 and 60 days after surgery in control (Fig. 2, , c). Newly formed cartilaginous tissue was formed only in the area of subchondral bone as a result of its injury, and it grew from the defect edges in the form of strip-like structures. In experiment the defect of articular cartilage was almost completely filled with newly formed hyaline cartilage having a smooth surface (Figure 2, b) which grew not only from the subchondral bone, but mainly from the defect edges where the isogenic groups of cells were located thereby, as evidenced by proliferative activity of chondrocytes (Fig. 2d).




Fig2: Zone of the defect of the rat knee cartilage 30 days after surgery: a, – control, the defect is filled with loose fibrillar connective tissue; b, d – experiment, the defect is filled with newly formed hyaline cartilage.(a, b - paraffin sections, staining with Alcian blue at 2.5, magnification 200; c - fibroblasts and collagen fibril, ultrathin sections, transmission electron microscopy, magnification 5000; d - chondrocytes are arranged as isogenic groups, scanning electron microscopy, magnification x 1300.)

The surface of newly formed hyaline cartilage acquired the shine characteristics of the articular cartilage of the intact animals. The cell population of the regenerated cartilage was mainly represented by proliferating chondrocytes; its structure did not possess the characteristics of the articular cartilage. Thirty days after surgery, the content of glycosaminoglycane non-sulphated and sulphated forms in the defect for experiment more than twice exceeded control values, but they did not differ from one another significantly. Sixty days after the surgery, glycosaminoglycane sulphated forms prevailed. It was evidenced by the higher maturity degree of the newly formed cartilaginous tissue (Table 1) which reached the maturity degree inherent for the articular cartilage of intact animals.

DISCUSSION

Mature hyaline cartilage is known to have a slight potential of restoration due to low density of cells and little mitotic activity of chondrocytes [9, 14]. Large defects of the articular cartilage are filled with biomechanically improper fibrillar cartilage followed by osteoarthritis development [10]. The advent of tissue engineering techniques has provided alternative possibilities to treat such patients by using cell therapy combined with synthetic substitutes of extracellular matrix and biologically active factors for functional replacement of articular hyaline cartilage [11, 12, 13]. The studies have demonstrated that the MBM granules when implanted into the defect zone of articular cartilage are identified in all the periods of further observation, they have marked chondromodulating effect, demonstrate prolonged activation of reparative chondro- and osteogenesis. Regenerated cartilage is formed early in the zone of articular cartilage injury, and it acquires the cell specificity of hyaline cartilage tissue. Integrated cartilage coating is formed, the disordered contours of articular surface are aligned gradually, thereby leading to complete or partial recovery of the joint functional activity. The newly formed cartilage dominated sulfated glycosaminoglycans, which indicates a high degree of maturity [14, 15]. Growth factors localized in the implanted MBM granules, as well as bone morphogenetic proteins being released during osteoclastic resorption provide chondro- and osteoinductor properties for the granules [16, 17, 18].

CONCLUSION

Thus, the studies have demonstrated that the use of the implant of granulated MBM as chondro- and osteogenesis stimulator and as a corrector of destructive disorders in bone and cartilaginous tissue in articular cartilage injuries seems theoretically justified and promising.

ACKNOWLEDGEMENT

We thank the staff of our institutions for their help in carrying out experiments and supervision over animals during all stages of work.

Conflict of Interest: Nil

REFERENCES

1. Deev RV, Isaev AA, Kochish AI, Tikhilov PM. Ways of development of cell technologies in bone surgery. *Travmatol Ortop Rossii*. 2008;47(1):65-74.
2. European Convention for the Protection of Vertebrate Animals used for Experimental and other Scientific Purposes. *Vopr Rekonstruktiv Plast Khirurgii*. 2003;(4):34-36.
3. Ir'ianov IuM, Ir'ianova TIu. Osteoplastic effectiveness of mineralized bone matrix. *Morfologiya*. 2013;143(1):63-68.
4. Ir'ianov IuM, Ir'ianova TIu. X-ray electron probe microanalysis in quantitative histochemistry. *Morfologicheskie vedomosti*. 2010;(3):77-81.
5. Lysenok LN. Biomaterials: contribution to the progress of modern medical technologies. *Kletochn Transplantol Tkan Inzheneriia*. 2005;(2):56-61.
6. Patent RF. Biomaterial for filling bone defects and a technique to obtain it. Ir'ianov IuM, Ir'ianova TIu. No 2478394. Filed 23.11.2011. *Publ. Bull* 2013;(10):66
7. Shishatskaia EI. Cell matrices of resorbable polyhydroxyalkanoates. *Kletochn Transplantol Tkan Inzheneriia*. 2007; 2(2):68-75.
8. Bessa PC, Casal M, Reis RL. Bone morphogenetic proteins in tissue engineering: the road from laboratory to clinic, part II (BMP delivery). *J Tissue Eng Regen Med*. 2008; 2(2-3):81-96.
9. Brown TD, Johnston RC, Saltzman CL, Marsh JL, Buckwalter JA. Posttraumatic osteoarthritis: a first estimate of incidence, prevalence, and burden of disease. *J Orthop Trauma*. 2006; 20(10):739-44.
10. Buckwalter JA. Articular cartilage injuries. *Clin Orthop Relat Res*. 2002; (402):21-37.
11. Danisovic L, Varga I, Zamborsky R, Böhmer D. The tissue engineering of articular cartilage: cells, scaffolds and stimulating factors. *Exp Biol Med (Maywood)*. 2012; 237(1):10-17.
12. Duguay N, Petite H, Arnaud . Biomaterials and osseous regeneration. *Ann Chir Plast Esthet*. 2000; 45():364-76.
13. Eberli D, Atala . Tissue engineering using adult stem cells. *Methods Enzymol*. 2006; 420:287-02.
14. Strehl R, Tallheden T, Sjögren-Jansson E, Minuth WW, Lindahl A. Long-term maintenance of human articular cartilage in culture for biomaterial testing. *Biomaterials*. 2005; 26(22):4540-45.
15. Chung C, Burdick JA. Influence of three-dimensional hyaluronic acid microenvironments on mesenchymal stem cell chondrogenesis. *Tissue Eng Part A* 2008;15:243–54
16. Bessa PC, Casal M, Reis RL. Bone morphogenetic proteins in tissue engineering: the road from laboratory to clinic, part II (BMP delivery). *J Tissue Eng Regen Med* 2008;2:81–96
17. Park KH, Na K. Effect of growth factors on chondrogenic differentiation of rabbit mesenchymal cells embedded in injectable hydrogels. *J Biosci Bioeng* 2008;106:74–79
18. Yamaoka H, Nishizawa S, Asawa Y, Fujihara Y, Ogasawara T, Yamaoka K, Nagata S, Takato T, Hoshi K. Involvement of fibroblast growth factor 18 in dedifferentiation of cultured human chondrocytes. *Cell Prolif* 2010; 43:67–76