

ISSN No: 2319-5886

International Journal of Medical Research & Health Sciences, 2017, 6(7): 67-69

# Pilomatrixoma in the Parotid Region - A Case Report

Meena Patil<sup>1</sup>\*, Ravi swami<sup>2</sup>

<sup>1</sup>Associate Professor, Bharati Vidyapeeth Dental College & Hospital, Pune, Maharashtra, India <sup>2</sup>Associate Professor, Bharati Vidyapeeth Medical College, Pune, Maharashtra, India

\*Corresponding e-mail: patilmeena64@gmail.com

# ABSTRACT

Pilomatrixoma is a relatively benign cutaneous adnexal neoplasm with differentiation towards the matrix and inner sheath of a normal hair follicle as well as hair cortex. It is commonly seen in head and neck region. We present a case of 49-year-old female presenting with a swelling in the preauricular region. She was referred for Fine Needle Aspiration Cytology (FNAC). On aspiration, smears showed picture of benign salivary gland neoplasm. However, histopathological diagnosis was pilomatrixoma.

Keywords: Fine needle aspiration cytology, Pilomatrixoma, Salivary gland

## INTRODUCTION

Pilomatrixoma, also known as calcifying epithelioma of Malherbe, is a slow-growing benign cutaneous adnexal neoplasm. It was first described by Malharbe and Chenantois in 1880. It was in 1961 that Forbes and Helwing suggested the name Pilomatrixoma [1,2].

Pilomatrixoma accounts for up to 0.2% of all routine dermatopathological specimens. The tumour occurs in all age groups. About 30% to 50% of cases present in young individuals less than 30 years of age. Previous studies have shown a female predominance [1].

Pilomatrixoma favour hair bearing areas with majority of cases arising in the head & neck region as well as upper extremities. Patient present with solitary, asymptomatic, slowly growing cystic or firm nodule 0.5-3 cm in diameter [1].

In this report, we present a case of pilomatrixoma localized in preauricular area (Parotid region).

# CASE REPORT

A 49-year-old female patient presented with complaints of swelling in the preauricular region. The swelling was a small painless nodule about  $1.5 \times 1$  cm slowly growing for 3 months. She was referred for fine needle aspiration cytology with clinical diagnosis suspicious of salivary gland neoplasm. Aspiration smears showed cohesive clusters of benign epithelial cells. Diagnosis offered was benign salivary gland neoplasm. The nodule was operated & biopsy sent for histopathological confirmation. On histopathology, sections showed densely cohesive basaloid cells & anucleate squamous cells (Ghost cells) in clusters along with multinucleate giant cells & areas of calcification (Figures 1-5). These features were suggestive of benign tumour of cutaneous appendages with differentiation towards hair follicle called as pilomatrixoma.



Figure 1 FNAC- Clusters of epithelial cells



Figure 2 FNAC- Clusters of epithelial cells



Figure 3 HP  $\times$  400- Ghost cells with giant cell



Figure 4 HP × 400- Ghost cells with giant cell



Figure 5 HP × 400- Ghost cells with calcification

#### DISCUSSION

Pilomatrixoma or Calcifying epithelioma of Malharbe, is a benign skin neoplasm that arises from hair follicle matrix cells. In 1880, Malharbe and Chenantais first described this lesion referred to as the calcifying epithelioma [3]. The term pilomatrixoma was introduced in a publication by Forbisand Helwing in 1961 to better convey the histologic course [4].

It may be seen in any age group, mostly occurring in children or in the first two decades of life. It usually arises in the head and neck region [5]. Pilomatrixoma clinically presents as a slow-growing, solitary, firm, well-circumscribed, painless dermal or subcutaneous tumour [6]. Histopathological examination is highly useful in the diagnosis of pilomatrixoma is difficult. Histopathological examination reveals round

islets of epithelial cells surrounded by a connective tissue capsule. In these islets, the eosinophilic ghost cells that have lost their nuclei are the specific indicators of pilomatrixoma. In addition to these islets, there are foreign body giant cells and basaloid cells with prominent nucleoli. Calcification is seen in 70-85% of the cases. The differential diagnosis of pilomatrixoma should include epidermoid cyst, foreign body granuloma, dermatofibroma, keratoacanthoma, nodular sub epidermal fibrosis, basal cell carcinoma, osteoma, and skin metastasis [7].

Likewise, on cytology, one can expect a varied cytomorphological spectrum depending on the stage of growth. Because of this varied presentation many authors have misdiagnosed pilomatrixomas on cytology. On cytological overview of literature, we found that pilomatrixomas on FNAC were diagnosed as malignant lesions [8,9]. Similarly, in our case, because of presence of cellular clusters of epithelial cells, absence of ghost cells, giant cells and calcification on cytology, location of lesion, we thought of benign salivary gland neoplasm of parotid gland; as these features better seen on histopathology.

Likewise, Luciano et al in his case report has mentioned that, pilomatrixoma was misdiagnosed on cytology as mucoepidermoid carcinoma because of the basaloid cells of pilomatrixoma were mistaken as intermediate cells of mucoepidermoid carcinoma [10]. Shivkumar, et al. also had diagnostic difficulty on fine needle aspiration which led to the three-differential diagnosis of mucoepidermoid carcinoma, squamous cell carcinoma and calcifying odontogenic tumour [11].

In this reported case, the definitive diagnosis was made after histologic examination following excision of the mass. Surgical excision is the method of choice in the treatment of pilomatrixoma since pilomatrixoma is benign lesion and dose not regress spontaneously. Postoperative recurrence of pilomatrixoma is extremely rare and occurs in 0-3% of the cases [7].

### CONCLUSION

Pilomatrixoma should be kept in mind in the differential diagnosis of head and neck tumours. The cytologic appearance can be misleading and hence excision with subsequent histopathologic examination will lead to the correct diagnosis.

### ACKNOWLEDGEMENT

We are very thankful to Dr. Shilpa Patel-Fernandis, V care Laboratory, Sinhgad Road, Pune for her support.

### REFERENCES

- [1] LeBoit, Philip E., ed. Pathology and Genetics of Skin Tumours. Vol. 10. IARC, 2006.
- [2] Brandner, Michael D., and Juris Bunkis. "Pilomatrixoma presenting as a parotid mass." *Plastic and Reconstructive Surgery* 78.4 (1986): 518-521.
- [3] Malherbe, A., and J. Chenantais. "Notes on calcifying epitheliomas of sebaceous glands." *Prog Med (Paris)* 8 (1880): 826-837.
- [4] Forbis, Robert, and Elson B. Helwig. "Pilomatrixoma (calcifying epithelioma)." Archives of Dermatology 83.4 (1961): 606-618.
- [5] Moehlenbeck, Friedrich W. "Pilomatrixoma (calcifying epithelioma): a statistical study." Archives of Dermatology 108.4 (1973): 532-534.
- [6] Wells, Neil J., et al. "Pilomatrixoma: A common, benign childhood skin tumour." Canadian Journal of Surgery 37.6 (1994): 483.
- [7] Agarwal, Ravi P., et al. "Pilomatrixoma of the head and neck in children." Otolaryngology—Head and Neck Surgery 125.5 (2001): 510-515.
- [8] Preethi, T. R., K. Jayasree, and E. K. Abraham. "A case of pilomatrixoma misdiagnosed as metastatic carcinoma on fine needle aspiration cytology." *Journal of Cytology* 24.4 (2007): 207.
- [9] Roche, N. A., S. J. Monstrey, and G. E. Matton. "Pilomatricoma in children: common but often misdiagnosed." Acta Chirurgica Belgica 110.2 (2010): 250-254.
- [10] MacDonald, Laurie, and Hossein M. Yazdi. "Non-diagnostic fine needle aspiration biopsy of the thyroid gland." Acta Cytologica 40.3 (1996): 423-428.
- [11] Sivakumar, Sivaselvam. "Pilomatrixoma as a diagnostic pitfall in fine needle aspiration cytology." *Acta Cytologica* 51.4 (2007): 583-585.