

- Stable metabolic status (i.e., electrolytes and glycemic index within normal range, body temperature $<38^{\circ}\text{C}$, Hemoglobinemia ≥ 8 g/dL-10 g/dL).
- Adequate oxygenation (i.e., arterial oxygen saturation (SaO₂) >92 % with inspiratory oxygen fraction (FiO₂) ≤ 0.5 or arterial oxygen partial pressure to inspiratory oxygen fraction (PaO₂/FiO₂) ≥ 150 mmHg, both with positive end-expiratory pressure (PEEP) ≤ 8 cmH₂O).
- Adequate pulmonary function (i.e., RR ≤ 30 breaths/min with VT ≥ 5 mL/kg Ideal Body Weight (IBW) and no significant respiratory acidosis).

Exclusion Criteria

- Inability to obtain proper acoustic windows.
- Inability to obtain written informed consent.
- Patients under 18 years old.
- Hemodynamically unstable patient, on vasopressors.
- Presence of thoracotomy, pneumothorax, or pneumomediastinum.
- Neuromuscular disease or use of muscle-paralyzing agents within 48 h before the study.
- Advanced Peripheral neuropathy patients.
- Presence of flail chest or rib fractures.
- Advanced malignancy.
- Advanced heart failure.

Diaphragmatic Functions

Each diaphragm was evaluated by B-mode and M-mode Subcostal views used to rule out any abnormalities in muscle movement. All measurements were done by the ultrasound probe of Siemens Acuson X300 PE ultrasound machine (probe of 2 mH-4 mH frequency). We assessed the following parameters.

Diaphragmatic Thickens Fraction

The diaphragm was visualized by placing the probe perpendicular to the chest wall, in the eighth or ninth intercostal space, between the anterior axillary and the midaxillary lines, to observe the zone of apposition of the muscle 0.5cm to 2 cm below the costophrenic sinus. The diaphragm was imaged as a structure with three distinct layers, including two parallel echoic lines (the diaphragmatic pleura and the peritoneal membrane) and a hypoechoic structure between them (the muscle itself) [16]. The patient was then instructed to perform full inspiration to Total Lung Capacity (TLC) and then to exhale to Residual Volume (RV). Several images of the diaphragm were then captured and stored, including at least three at the point of maximum thickening at TLC and at least three at minimum thickening at RV. Images were obtained within the first 5 min of the SBT to identify the diaphragm and measure its thickness. It took three to five sessions; 10 min-15 min each.

On each frozen B-mode image; the diaphragm thickness was measured from the middle of the pleural line to the middle of the peritoneal line [16].

Then; DTF was calculated as a percentage from the following formula: Thickness at end inspiration-Thickness at end-expiration/Thickness at end-expiration (Figure 1) [17-23].



Figure 1. Ultrasound B-mode using 7.5MHz probe in the zone of apposition. The diaphragm muscle appeared as hypochoic structure between the diaphragmatic pleura and peritoneal pleura (the two echoic lines) (arrow).




Figure 1 (A,B) Diaphragmatic thickens fraction

Diaphragmatic Excursion

- At the Anterior axillary line in the subcostal region, the Transducer is directed medially, cranially, and dorsally.
- B mode was used to assess excursion and M mode to measure the Normal range of motion from the resting expiratory position to full inspiration in adults 1.9 cm to 9 cm.
- Excursion >2.5 cm in adults as a cut-off was used for excluding severe diaphragm dysfunction [5].

Diaphragmatic Dysfunction

Diaphragmatic dysfunction was defined, during quiet breathing as Diaphragmatic excursion ≤ 11 mm or less and Diaphragm thickness less than 0.2 cm [5]. The Diaphragmatic Rapid Shallow Breathing Index (D-RSBI).

Statistical Analysis

Numerical variables were described as mean \pm SD. Categorical variables were described as percentages. Comparisons were done using Student "t" and one-way ANOVA tests for numerical variables and the Chi-square test for categorical variables. Correlations were plotted and r values (correlation coefficients) were stated. ROC curves were plotted to determine cut-offs. P value was considered significant if ≤ 0.05 . Statistics were calculated using SPSS 21 package [17].

RESULTS

In our study, DTF (Diaphragmatic Thickening Fraction), DE (Diaphragmatic Excursion), RSBI (Rapid Shallow Breathing Index), and D-RSBI (Diaphragmatic Rapid Shallow Breathing Index) was calculated for all those patients and correlated with the possibility of weaning. Weaned patients from mechanical ventilation were also followed up within the upcoming 48 hours for the risk of reintubation.

The study population was subdivided into two groups; 26 patients (65%) were weaned successfully and 14 patients (35%) failed the weaning trial (Table 1 and 2).

Table 1 Shows the study's descriptive data concerning the baseline

		Range	Mean ± SD
Age	Failure	56-91	71.2 ± 10.7
	Success	22-75	4 ± 14.4
ICU stay (days)	Failure	9-45	26.5 ± 10.
	Success	7-60	18.8 ± 12.9
Ventilation period	Failure	3-14	6.85 ± 3.65
	Success	1-16	5.46 ± 4.26
Ventilatory parameters			
PaO₂/FiO₂	Failure	144-306	253.4 ± 50
	Success	234-350	298.3 ± 26.7
RSBI(breaths/min/L)	Failure	45-90	76.5 ± 15.8
	Success	33-8-	52 ± 12.9
C stat (ml/cmH₂O)	Failure	35-80	53.7 ± 11.4
	Success	50-77	63.3 ± 6.8
Resistance(cmH₂O/L/sec)	Failure	3-15	6.9 ± 2.8
	Success	14-19	11.3 ± 4.5
VT(ml)	Failure	280-430	330.7 ± 36.2
	Success	350-600	437.5 ± 57.6
P peak (cmH₂O)	Failure	21-34	28.3 ± 3.5
	Success	21-32	24.2 ± 2.9
P plateau (cmH₂O)	Failure	14-22	17.01 ± 2.8
	Success	14-23	17.3 ± 1.78
Chest U/S parameters			
DTF	Failure	0.11-0.73	0.31 ± 0.21
	Success	0.29-0.96	0.56 ± 0.22
Excursion(mm)	Failure	7-16	12 ± 3
	Success	16-32	21 ± 4
DRBSI (breaths/min/mm)	Failure	2.7 ± 0.9	1.4-4.2
	Success	1.37 ± 0.46	0.8-2.6
Scores			
SOFA	Failure	6.7 ± 2.4	03-11
	Success	3.6 ± 3.03	0-11

APACHE II	Failure	17.2 ± 4.2	12-24
	Success	9.8 ± 4.5	4-17
Expected Mortality	Failure	24.2 ± 11.	12-40
	Success	9 9.8 ± 7.3	1-26

Table 2 Shows the study's descriptive data ON successful and unsuccessful weaning

Demographic data	Weaning failure	Weaning success
Sex		
Male	10 (40%)	15 (60%)
Female	4 (26.6%)	11 (74.4%)
Comorbidities		
HTN	3 (27%)	8 (73%)
DM	5 (45%)	6 (55%)
Heart failure	6 (33%)	12 (67%)
Pneumonia	13 (46%)	15 (54%)
Renal impairment	10 (62.5%)	6 (37.5%)
Hypothyroidism	4 (50%)	4 (50%)
Neurological disorders	6 (55%)	5 (45%)
Fate		
Survivor	8 (57%)	23 (88.5%)
Non-survivor	6 (43%)	3 (11.5%)

Analytical Data

Shows the relation between age, gender, and weaning results where there was a strong statistical relationship between age and weaning failure with a p-value of 0.001 (Table 3).

Table 3 Relation between age, gender, and weaning results

	Weaning		p-value
Gender	Failure	Successful	0.307
Male	4 (26.6%)	11 (74.4%)	
female	10 (40%)	15 (60%)	
Age	Mean ± SD	0.001	
Failure	71.2 ± 10.7		
Successful	55.4 ± 14.4		

Relation between different risk factors and comorbidities and weaning outcomes where pneumonia and renal impairment show a strong association with weaning failure with a p-value of 0.021 and 0.004 respectively (Table 4).

Table 4 Relation between different risk factors and comorbidities

Comorbidities	Weaning		p-value
	Failure	Successful	
HTN	3 (27%)	8 (73%)	0.404

DM	5 (45%)	6 (55%)	0.311
Heart failure	6 (33%)	12 (67%)	0.554
Pneumonia	13 (46%)	15 (54%)	0.021
Renal impairment	10 (62.5%)	6 (37.5%)	0.004
Hypothyroidism	(50%) 4	4 (50%)	0.276
Neuro disorders	6 (55%)	5 (45%)	0.111

The relation between risk factors and mortality where pneumonia shows a strong association with mortality with a p-value of 0.016 (Table 5).

Table 5 Relation between risk factors and mortality with a p-value of 0.016

Risk factors	Mortality		p-value
	survivor	Nonsurvivor	
DM	9 (81%)	2 (19%)	0.523
Pneumonia	5 (18%)	21 (82%)	0.016
HTN	9 (81%)	2 (19%)	0.523
Heart failure	15 (83.4%)	3 (16.6%)	0.341
Neuro disorder	8 (73%)	3 (27%)	0.477
Hypothyroidism	5 (62.5%)	3 (37.5%)	0.477
Renal impairment	10 (62.5%)	6 (37.5%)	0.042

The relation between ventilator parameters and weaning outcome where all mentioned parameters show a strong association with weaning outcome (Table 6).

Table 6 Relation between ventilator parameters and weaning outcome

Mean ± SD			p-value
Ventilator measurements			
	Weaning Failure	Weaning Successful	
PaO ₂ /FiO ₂	253.4 ± 50	298.3 ± 26.7	0.006
RSBI	67.5 ± 15.8	52 ± 12.9	0.008
Static compliance	53.7 ± 11.4	63.3 ± 6.8	0.002
Resistance	11.3 ± 4.5	6.9 ± 2.8	0.004

The relation between LUS measurements and weaning outcome, where all measurements show a strong association with weaning outcome (Table 7).

Table 7 Relation between LUS measurements and weaning outcome

	Mean ± SD		p-value
	Weaning Failure	Weaning Successful	
Thickness fraction	0.31 ± 0.21	0.56 ± 0.22	0.001
excursion	1.24 ± 0.3	2.11 ± 0.4	<0.001
DRBSI	2.7 ± 0.9	1.37 ± 0.46	<0.001

This shows the relation between weaning outcomes sepsis, and mortality scores where both SOFA score and APACHE II score show a strong association with weaning outcome with p-value 0.021 and 0.032 respectively (Table 8).

Table 8 Relation between weaning outcomes sepsis, and mortality scores where both SOFA score and APACHE II

Scores	Mean \pm SD		p-value
	Weaning Failure	Weaning Successful	
SOFA	6.7 \pm 2.4	3.7 \pm 3.0	0.021
APACHE II	17.2 \pm 4.2	9.8 \pm 4.5	0.032

Correlative Data

The Correlations of age, DTF, and DE with D-RSBI and RSBI where age and DE show strong statistical association with p-values 0.01 and <0.001 respectively, and r-values 0.353 and 0.729 respectively (Table 9).

Table 9 Statistical association with p-values 0.01 and <0.001 respectively and r values 0.353 and 0.729 respectively

Variables		D-RSBI	RSBI
Age	Pearson Correlation(r)	0.353	0.463
	p-value	0.025	0.002
DTF	Pearson Correlation(r)	0.401	0.221
	p-value	0.01	0.169
DE	Pearson Correlation(r)	0.729	0.478
	p-value	<0.001	0.002

Shows the Correlations of P/F ratio, RSBI and DTF with the length of ICU stay where statistical association showed p-value 0.004,0.016 and 0.013 respectively and r-value 0.437,0.377 and 0.389 respectively (Table 10).

Table 10 Correlations of P/F ratio, RSBI, and DTF with the length of ICU

Variables		Length of ICU stay
P\F ratio	Pearson Correlation (r)	0.437
	p-value	0.004
RSBI	Pearson Correlation (r)	0.377
	p-value	0.016
DTF	Pearson Correlation (r)	0.389
	p-value	0.013

Comparative Data

Comparative statistical analysis between different outcomes including weaning results and mortality showed a significant statistical association with a p-value of 0.033 (Table 11).

Table 11 Comparative statistical analysis between different outcomes

fate	Weaning		p-value
	Failure (14)	Successful (26)	
survivors(31)	8 (57%)	23 (88.5%)	0.033
Non survivors(9)	6 (43%)	3 (11.5%)	

Comparison between survivors and non-survivors as regards chest U/S where Thickness fraction, Excursion, and D-RSBI showed a p-value of 0.004, 0.005, and 0.036 respectively (Table 12).

Table 12 Comparison between survivors and non-survivors

LUS measurements	Survivor	Non-survivor	p-value
	Mean \pm SD		
Thickness fraction	0.5 \pm 0.3	0.3 \pm 0.1	0.004
Excursion	1.9 \pm 0.6	1.7 \pm 0.5	0.005
D-RSBI	2.1 \pm 2.2	1.0 \pm 1.0	0.036

Receiver Operating Characteristic (ROC) Curve Analysis

For prediction of successful weaning by the use of LUS parameters, RSBI, DSBI. At a cut of 30% of DTF, its sensitivity to predict successful weaning was 96% and its specificity was 71 % Area Under the ROC Curve (AUC=0.676). At a cut of 17 mm of diaphragmatic Excursion, its sensitivity to predict successful weaning was 88% and its specificity was 100% (AUC=0.973). At a cut of 61 breaths/min/L of RSBI, its sensitivity to predict successful weaning was 73% and its specificity was 71 % (AUC=0.755). At a cut of 1.8 breaths/min/mm of D-RSBI, its sensitivity to predict successful weaning was 84% and its specificity was 92% (AUC=0.935) (Figure 2 and 3).




Figure 2 ROC curve for prediction of successful weaning by RSBI and D-SBI




Figure 2 ROC curve for prediction of successful weaning by LUS

DISCUSSION

The diaphragmatic Rapid Shallow Breathing Index (D-RSBI), the ratio between respiratory rate (RR) and the ultrasonographic Diaphragmatic Displacement (DD), is a new and promising tool to predict weaning outcomes [18]. Our results showed a statistically significant relationship between age and weaning results with a p-value (of 0.001). In contrast to our study, there was no statistically significant relationship between age and weaning results in Spadaro et al., 2016, Walaa and Lutfy., 2018 and Mowafy and Abdelgalel, 2018, with p-value (0.338, 0.508, and 0.976) respectively [19]. This can be explained by different age population groups, our study groups' mean age was 55.4 and SD 14.4 for successful weaning groups, and the mean age was 71.2 and SD 10.7, while for Spadaro et al., 2016 the study groups' mean age was 62 and SD 16 for successful weaning groups and mean age 66 and SD 11, relatively younger age groups for Walaa and Lutfy, 2018 and Mowafy and Abdelgalel, 2018.

There were no statistically significant differences between comorbidities and weaning outcomes in the two study groups except for pneumonia and renal impairment with p-values of 0.021 and 0.004 respectively. In the same context, Spadaro et al., 2016 and (Mowafy and Abdelgalel, 2018) did not show significant statistical relation between weaning outcome and comorbidities except for sepsis as a reason for initiating mechanical ventilation ($p=0.035$ and 0.008) respectively. This can be explained by different age groups, comorbidities, and types of patients as our study was conducted on medical and surgical ICU patients while for Spadaro et al., 2016, the study was conducted on medical ICU patients only and for Mowafy and Abdelgalel, 2018 the study was conducted on COPD patients and severity of illness [15,20,21].

PaO_2/FiO_2 ratio was calculated before the start of SBT to indicate readiness for the weaning process, and then it was correlated with different variables. PaO_2/FiO_2 ratio was found to be significantly lower for patients who failed to wean than for those who were successfully weaned ($p=0.006$). The P/F ratio of the successful weaning groups ranged from 234 to 350 with a mean value and standard deviation (of 298.3 ± 26.7) while the failed weaning groups ranged from 144 to 306 with a mean value and standard deviation (of 253.4 ± 50). In concordance with our study, PaO_2/FiO_2 ratio for Roche-Campo et al., 2018 and Moschietto et al., 2012 was found to be significantly higher for patients who successfully weaned from MV ($p \leq 0.001$) ($p=0.002$) respectively [22,23].

Regarding weaning from mechanical ventilation, our results showed that the cut-off point of DTF for successful weaning from mechanical ventilation was $\geq 30\%$. In the same context, the other cut-off points were in DiNino et al., 2014 $\geq 30\%$, Ferrari et al., 2014 >36 , Shereen et al., 2016 $\geq 34.2\%$, Agmy et al., 2015 $>40\%$ and Soliman et al., 2019 $>30\%$ and all were associated with a successful SBT [24-27]. In our results, the DTF of the successful weaning groups ranged from 0.29 to 0.96 with a mean value of 0.56 while for the failed weaning groups it ranged from 0.11 to 0.73 with a mean value of 0.3. In the same context, in Ferrari et al., 2014 the DTF of the successful weaning groups ranged from 0.38 to 0.64 with a mean value of 0.56 while the failed weaning groups ranged from 0.22 to 0.30 with a mean value of 0.26. In our results, the DTF of the successful weaning groups was found to be significantly higher than the failed groups from MV ($p=0.001$). In concordance with our study, in Ferrari et al., 2014, (Soliman et al., 2019), and Shereen et al., 2016, the DTF of the successful weaning groups was found to be significantly higher than in the failed groups from MV ($p \leq 0.001$) ($p=0.001$) ($p<0.05$) respectively [28]. Differences in DTF cut-off values can be explained by differences in methodology (DiNino et al., 2014), the measurements were taken 12 hours to 36 hours before extubation, and Shereen et al., 2016 were done in controlled and spontaneous mode, variable inspiratory efforts, the patient population as in (Agmy et al., 2015 the study was carried on COPD patients, the severity of illness at ICU admission, metabolic conditions, and duration of MV (Soliman et al., 2019) and Ferrari et al., 2014 were all patients on tracheostomy tube [25,28].

In our study, the cut-off point of DE that can predict successful weaning was ≥ 17 mm at end of inspiration. This was close to the cut-off point of Shereen et al., 2016 (44) which was >11 mm, Osman et al., 2017 >22 mm, Kim et al., 2011 >17.9 mm, Spadaro et al., 2016 ≥ 14 mm and Walaa and Lutfy, 2018 >12 mm [15,18,29]. Our results showed that DE was significantly higher (21 mm (16 mm-32 mm) *vs* 12 mm (7 mm-16 mm) respectively in the successful extubation group compared to the failed one. In concordance with our study, Shereen et al., 2016 stated that DE was significantly higher (16 mm (11.4 mm-18 mm) *vs* 9.8 mm (8.8 mm-12 mm) respectively) in the successful extubation group compared to the failed one. In our results, the DE of the successful weaning groups was found to be significantly higher than the failed groups from MV ($p \leq 0.001$). In the same context, in Spadaro et al., 2016 and Walaa and Lutfy, 2018, the DE of the successful weaning groups was found to be significantly higher than the failed groups from MV ($p \leq 0.001$) ($p \leq 0.001$) respectively. Different results of DE may be due to differences in posture, showing higher values when patients are supine versus seated, as well as when the abdominal and/or thoracic pressure is altered (eg. ascites, atelectasis) Kim et al., 2011, patient population (Walaa and Lutfy, 2018 were carried on COPD patients, Osman et al., 2017 was carried on postoperative patients, in Spadaro et al., 2016 spontaneous ventilation through a T-tube was done with the FiO_2 set at the same level used during mechanical ventilation, Also diaphragmatic ultrasound requires the use of high-frequency probes with precision to a few millimeters, meaning that slight variations in measurement between observers can substantially affect the result [30-33].

Our results showed that the cut-off point of RSBI for successful weaning from mechanical ventilation was ≤ 61 breaths/min/L showed 73% sensitivity and 71% specificity (AUC=0.755) was associated with a successful SBT. In the same context, the other cut-off points were Spadaro et al., 2016 ≥ 62 breaths/min/L showed 52% sensitivity and 97% specificity (AUC=0.720). , Walaa and Lutfy, 2018 (18) <70 breaths/min/L showed 69% sensitivity and 71%

specificity(AUC=0.669), and (Soliman et al., 2019) <71 breaths/min/L showed 80% sensitivity and 70% specificity(AUC=0.815)., and Pirompanich and Romsaiyut, 2018 <105 breaths/min/L showed 78% sensitivity and 92% specificity and all were associated with a successful SBT. Our results showed that RSBI was significantly lower [52 breaths/min/L (33-80) vs 67 breaths/min/L (45-90) respectively] in the successful extubation group compared to the failed one [34]. In the same context Spadaro et al., 2016 stated that RSBI was significantly lower breaths/min/L (32-52) vs 63 breaths/min/L (37-90) respectively] in the successful extubation group compared to the failed one. In our results, the RSBI of the successful weaning groups was found to be significantly lower than the failed groups from MV (p=0.011). In concordance with our study, in Spadaro et al., 2016, the RSBI of the successful weaning groups was found to be significantly lower than the failed groups from MV (p=0.012) while in concordance with our study, in Walaa and Lutfy, 2018, there was no statistically significant difference between failed and successful weaning groups (p=0.082) [18]. In concordance with our study, patients who failed the weaning attempt in this study had a lower RSBI (75.5 ± 11.4 breaths/min/L) than the cutoff value of 105 breaths/min/L to predict weaning failure originally described by Yang and Tobin [35]. Several other studies have documented a wide range of predictive values for RSBI, which could reflect Differences in methodology (DiNino et al, the measurements were taken 12-36 hours before extubation, Pirompanich and Romsaiyut, 2018 SBT was done for 1 hour and Spadaro et al., 2016 was done on medical ICU patients), variable inspiratory efforts, patient population Walaa and Lutfy, 2018 was carried on COPD patients, the severity of illness at ICU admission (Soliman et al., 2019) [15,18,36].

Our results showed that the cut-off point of D-RSBI for successful weaning from mechanical ventilation was ≤ 1.8 breaths/min/mm showed 84% sensitivity and 92% specificity (AUC=0.935) was associated with a successful SBT. In the same context, the other cut-off points were Spadaro et al., 2016 (15)< 1.3 breaths/min/mm yielded 94.1 % sensitivity, 64.7% specificity (AUC=0.890), Walaa and Lutfy, 2018 <1.9 showed 82% sensitivity and 93% specificity (AUC=0.970) and (Mowafy and Abdelgalel, 2018) <1.6 breaths/min/mm showed 84% sensitivity and 100% specificity (AUC=0.979) and all were associated with a successful SBT [19]. Our results showed that D-RSBI was significantly lower (1.3 breaths/min/mm (0.8-2.6) vs 2.5 breaths/min/mm (1.4-4.2) respectively) in the successful extubation group compared to the failed one. In the same context Spadaro et al., 2016 stated that D-RSBI was significantly lower [1.2 breaths/min/mm (0.8-1.9) vs 2.7 breaths/min/mm (1.7-4.1) respectively] in the successful extubation group compared to the failed one. In our results, the D-RSBI of the successful weaning groups was found to be significantly lower than the failed groups from MV (p \leq 0.001). In the same context, in Spadaro et al., 2016 and Walaa and Lutfy, 2018, the D-RSBI of the successful weaning groups was found to be significantly lower than the failed groups from MV (p \leq 0.001) (p \leq 0.001) respectively. Different results of D-RSBI can be explained by differences in posture, showing lower values when patients are supine versus seated, as well as methodology, patient population(Walaa and Lutfy, 2018 were carried on COPD patients, in Spadaro et al., 2016 spontaneous ventilation through a T-tube was done requiring more effort than SBT and (Mowafy and Abdelgalel, 2018) was done on surgical ICU patients). Also, diaphragmatic ultrasound variations in measurement between observers can affect the result. There is a strong positive correlation between D-RSBI and RSBI (r=0.678). Also, a statistically significant difference was found between DRSBI and RSBI (p \leq 0.001), similar to Spadaro et al., 2016

($p \leq 0.001$). A systematic review meta-analysis held a comparison between DTF and DE and found that DE is associated with inspiratory volume but does not correlate with other indexes of respiratory effort and data suggest a lower accuracy for DE compared to DTF in predicting weaning outcome [37]. Moreover, in patients undergoing MV, DTF reflects active diaphragm contraction, while DE comes from adding patients' respiratory effort to the pressure generated by the ventilator, so its use is only meaningful in the absence of ventilatory support [37]. Despite being an observer-dependent technique, the available evidence suggests that both DTF and DE are reproducible measures and this can explain the difference in our results, especially in a diaphragmatic excursion to other studies. The D-RSBI was speculated to be more accurate than the RSBI because DD reflects more closely the diaphragmatic function as compared to VT. In the presence of diaphragmatic dysfunction, the diaphragm movement is depressed and the accessory muscles of inspiration assume a greater role in generating VT, but for a limited period as they are less efficient and easily fatigable [38,39]. In these situations, the VT is less linked to the function of the diaphragm than DD, which reflects the ability of the diaphragm to generate inspiratory volume [40]. However, while careful clinical observation remains the most important step in the decision to extubate, integrating it with the result of a relatively easy-to-obtain index like the D-RSBI could at least support the clinician in their decision [41-44].

LIMITATIONS

This study could be summarized in:

- A single physician performed the US.
- The nutritional status of the patients was not assessed which may affect muscle power.
- Being a single-center study.

CONCLUSION

DD(defined as $DD < 10$ mm), when combined with RR in an index that we named D-RSBI (RR/DD), is more accurate than the traditional RSBI (RR/VT) in predicting the weaning outcome, where, D-RSBI (RR/DD) cut-off of 1.8 is associated with the best sensitivity and specificity. We confirmed the importance of rapid and shallow breathing as a global index of weaning-induced patient distress.

DECLARATIONS

Conflict of Interest

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

REFERENCES

- [1] Pinsky, Michael R., and Jean-Francois Dhainaut, eds. "Pathophysiologic foundations of critical care." Williams and Wilkins, 1993.
- [2] Esteban, Andrés, et al. "Extubation outcome after spontaneous breathing trials with T-tube or pressure support ventilation." *American journal of respiratory and critical care medicine*, Vol. 156, No. 2, 1997, pp. 459-65.
- [3] Vassilakopoulos, T., S. Zakynthinos, and C. H. Roussos. "Respiratory muscles and weaning failure." *European Respiratory Journal*, Vol. 9, No. 11, 1996, pp. 2383-2400.

- [4] Heunks, Leo M., and Johannes G. Van Der Hoeven. "Clinical review: The ABC of weaning failure-a structured approach." *Critical care*, Vol. 14, No. 6, 2010, pp. 1-9.
- [5] McCool, F. Dennis, and George E. Tzelepis. "Dysfunction of the diaphragm." *New England Journal of Medicine*, Vol. 366, No. 10, 2012, pp. 932-42.
- [6] Jaber, Samir, et al. "Rapidly progressive diaphragmatic weakness and injury during mechanical ventilation in humans." *American journal of respiratory and critical care medicine*, Vol. 183, No. 3, 2011, pp. 364-71.
- [7] Berek, K., et al. "Polyneuropathies in critically ill patients: a prospective evaluation." *Intensive care medicine*, Vol. 22, No. 9, 1996, pp. 849-55.
- [8] Coakley, J. H., et al. "Preliminary observations on the neuromuscular abnormalities in patients with organ failure and sepsis." *Intensive care medicine*, Vol. 19, No. 6, 1993, pp. 323-28.
- [9] Coakley, J. H., et al. "Preliminary observations on the neuromuscular abnormalities in patients with organ failure and sepsis." *Intensive care medicine*, Vol. 19, No. 6, 1993, pp. 323-28.
- [10] Kim, W. Young, et al. "Diaphragm dysfunction assessed by ultrasonography: influence on weaning from mechanical ventilation." *Critical care medicine*, Vol. 39, No. 12, 2011, pp. 2627-30.
- [11] Yang, Karl L., and Martin J. Tobin. "A prospective study of indexes predicting the outcome of trials of weaning from mechanical ventilation." *New England Journal of Medicine*, Vol. 324, No. 21, 1991, pp. 1445-50.
- [12] Lee, Kang H., et al. "Rapid shallow breathing (frequency-tidal volume ratio) did not predict extubation outcome." *Chest*, Vol. 105, No. 2, 1994, pp. 540-43.
- [13] Savi, Augusto, et al. "Weaning predictors do not predict extubation failure in simple-to-wean patients." *Journal of critical care*, Vol. 27, No. 2, 2012, pp. 221-e1.
- [14] Huang, Chun-Ta, and Chong-Jen Yu. "Conventional weaning parameters do not predict extubation outcome in intubated subjects requiring prolonged mechanical ventilation." *Respiratory care*, Vol. 58, No. 8, 2013, pp. 1307-14.
- [15] Spadaro, Savino, et al. "Can diaphragmatic ultrasonography performed during the T-tube trial predict weaning failure? The role of diaphragmatic rapid shallow breathing index." *Critical Care*, Vol. 20, No. 1, 2016, pp. 1-11.
- [16] Wait, J. L., and R. L. Johnson. "Patterns of shortening and thickening of the human diaphragm." *Journal of Applied Physiology*, 1997.
- [17] Statistics, I. S. "IBM Corp. Released 2013. IBM SPSS Statistics for Windows, Version 22.0. Armonk, NY: IBM Corp." *Google Search*, 2013.
- [18] Meghawri, Mohammed Walaa Saeed. "Role of diaphragmatic rapid shallow breathing index in predicting weaning outcome in patients with acute exacerbation of COPD." *International Journal of COPD*, 2018.
- [19] Mowafy, Sherif MS, and Essam F. Abdelgalel. "Diaphragmatic rapid shallow breathing index for predicting weaning outcome from mechanical ventilation: Comparison with traditional rapid shallow breathing index." *Egyptian Journal of Anaesthesia*, 2018.
- [20] DiNino, Ernest, et al. "Diaphragm ultrasound as a predictor of successful extubation from mechanical ventilation." *Thorax*, Vol. 69, No. 5, 2014, pp. 431-435.

- [21] Ferrari, Giovanni, et al. "Diaphragm ultrasound as a new index of discontinuation from mechanical ventilation." *Critical ultrasound journal*, Vol. 6, No. 1, 2014, pp. 1-6.
- [22] Moschietto, Sebastien, et al. "Transthoracic echocardiography with Doppler tissue imaging predicts weaning failure from mechanical ventilation: evolution of the left ventricle relaxation rate during a spontaneous breathing trial is the key factor in weaning outcome." *Critical care*, Vol. 16, No. 3, 2012, pp. 1-10.
- [23] DiNino, Ernest, et al. "Diaphragm ultrasound as a predictor of successful extubation from mechanical ventilation." *Thorax*, Vol. 69, No. 5, 2014, pp. 431-35.
- [24] Ferrari, Giovanni, et al. "Diaphragm ultrasound as a new index of discontinuation from mechanical ventilation." *Critical ultrasound journal*, Vol. 6, No. 1, 2014, pp. 1-6.
- [25] Ferrari, Giovanni, et al. "Diaphragm ultrasound as a new index of discontinuation from mechanical ventilation." *Critical ultrasound journal*, Vol. 6, No. 1, 2014, pp. 1-6.
- [26] Farghaly, Shereen, and Ali A. Hasan. "Diaphragm ultrasound as a new method to predict extubation outcome in mechanically ventilated patients." *Australian Critical Care*, Vol. 30, No. 1, 2017, pp. 37-43.
- [27] Agmy, Gamal, Samiaa Hamdy, and Sherin Farghally. "Diaphragm ultrasound as a novel guide of weaning from invasive ventilation." 2015.
- [28] Soliman, Soliman Belal, et al. "Chest ultrasound in predication of weaning failure." *Open access Macedonian journal of medical sciences*, Vol. 7, No. 7, 2019, p. 1143.
- [29] Osman, Ahmed M., and Reham M. Hashim. "Diaphragmatic and lung ultrasound application as new predictive indices for the weaning process in ICU patients." *The Egyptian Journal of Radiology and Nuclear Medicine*, Vol. 48, No. 1, 2017, pp. 61-66.
- [30] Ambrosino, Nicolino, and Luciano Gabbrielli. "The difficult-to-wean patient." *Expert review of respiratory medicine*, Vol. 4, No. 5, 2010, pp. 685-92.
- [31] Matamis, Dimitrios, et al. "Sonographic evaluation of the diaphragm in critically ill patients. Technique and clinical applications." *Intensive care medicine*, Vol. 39, No. 5, 2013, pp. 801-10.
- [32] Boussuges, Alain, Yoann Gole, and Philippe Blanc. "Diaphragmatic motion studied by m-mode ultrasonography: methods, reproducibility, and normal values." *Chest*, Vol. 135, No. 2, 2009, pp. 391-400.
- [33] Wait, Juliette L., et al. "Diaphragmatic thickness-lung volume relationship in vivo." *Journal of Applied Physiology*, Vol. 67, No. 4, 1989, pp. 1560-68.
- [34] Pirompnich, Pattarin, and Sasithon Romsaiyut. "Use of diaphragm thickening fraction combined with rapid shallow breathing index for predicting success of weaning from mechanical ventilator in medical patients." *Journal of intensive care*, Vol. 6, No. 1, 2018, pp. 1-7.
- [35] Yang, Karl L., and Martin J. Tobin. "A prospective study of indexes predicting the outcome of trials of weaning from mechanical ventilation." *New England Journal of Medicine*, Vol. 324, No. 21, 1991, pp. 1445-50.
- [36] DiNino, Ernest, et al. "Diaphragm ultrasound as a predictor of successful extubation from mechanical ventilation." *Thorax*, Vol. 69, No. 5, 2014, pp. 431-35.
- [37] Llamas-Alvarez, Ana M., Eva M. Tenza-Lozano, and Jaime Latour-Perez. "Diaphragm and lung ultrasound to predict weaning outcome: systematic review and meta-analysis." *Chest*, Vol. 152, No. 6, 2017, pp. 1140-50.

- [38] Yan, S. H. E. N. G., et al. "Effect of diaphragmatic fatigue on control of respiratory muscles and ventilation during CO₂ rebreathing." *Journal of Applied Physiology*, Vol. 75, No. 3, 1993, pp. 1364-70.
- [39] Yan, S. H. E. N. G., et al. "Effect of global inspiratory muscle fatigue on ventilatory and respiratory muscle responses to CO₂." *Journal of Applied Physiology*, Vol. 75, No. 3, 1993, pp. 1371-77.
- [40] Umbrello, Michele, et al. "Diaphragm ultrasound as indicator of respiratory effort in critically ill patients undergoing assisted mechanical ventilation: a pilot clinical study." *Critical Care*, Vol. 19, No. 1, 2015, pp. 1-10.
- [41] Esteban, Andrés, et al. "A comparison of four methods of weaning patients from mechanical ventilation." *New England Journal of Medicine*, Vol. 332, No. 6, 1995, pp. 345-50.
- [42] Brochard, Laurent, et al. "Comparison of three methods of gradual withdrawal from ventilatory support during weaning from mechanical ventilation." *American journal of respiratory and critical care medicine*, Vol. 150, No. 4, 1994, pp. 896-03.
- [43] Alvisi, R., et al. "Predictors of weaning outcome in chronic obstructive pulmonary disease patients." *European Respiratory Journal*, Vol. 15, No. 4, 2000, pp. 656-62.
- [44] Farghaly, Shereen, and Ali A. Hasan. "Diaphragm ultrasound as a new method to predict extubation outcome in mechanically ventilated patients." *Australian Critical Care*, Vol. 30, No. 1, 2017, pp. 37-43.