



Prevalence of Social Anxiety Disorder and Its Impact on Academic Performance among Female Secondary School Students in Saudi Arabia

Tharwat Abdulaziz Alahmary^{1*}, Riyadh A. Algamdi¹, Mohammad Abdulhameed Alharbi² and Sami Abdo Radman Al-Dubai³

¹Joint Program of Family Medicine Post Graduate Studies, Ministry of Health, Al-Madinah, Saudi Arabia

²Faculty of Medicine, King Abdulaziz University, Jeddah, Saudi Arabia

³Joint Program of Preventive Medicine Post Graduate Studies, Ministry of Health, Al-Madinah, Saudi Arabia

*Corresponding e-mail: tharwat.alahmary@gmail.com

Received: 29-September-2022, Manuscript No. ijmrhs-22-66252; **Editor assigned:** 03-October-2022, PreQC No. ijmrhs-22-66252 (PQ); **Reviewed:** 18-October-2022, QC No. ijmrhs-22-66252 (Q); **Revised:** 25-October-2022, Manuscript No. ijmrhs-22-66252 (R); **Published:** 30-October-2022, J-invoice: J-66252

ABSTRACT

Background: Social Anxiety Disorder (SAD) is a common type of anxiety disorder among females that impacts their quality of life and academic performance if not discovered and treated early. **Objectives:** To determine the prevalence, severity, and associated factors of SAD and to assess its impact on the academic performance of female secondary school students in Medina. **Subjects and methods:** This cross-sectional study included a representative sample of Saudi girls enrolled in governmental secondary schools in Al-Madinah. A self-administered questionnaire was used for data collection. It included socio-demographic data, the Social Phobia Inventory (SPIN) to detect SAD, and the Leibowitz Social Anxiety Scale (LSAS) to evaluate the severity of SAD. **Results:** The study included 405 female students. The majority were aged ≤ 17 years (58.2%). The prevalence of SAD was 51.1%. Marked, severe, and very severe SAD was reported by 12.8%, 4.7%, and 3.7% respectively. SAD was higher among first birth order students, those with less-educated parents, and those with lower income ($p < 0.05$). SAD was significantly associated with all aspects of the paternal-students relationship ($p < 0.05$). **Conclusion:** SAD is common among girls in secondary schools in Medina. SAD was significantly associated with income, level of parental education, and paternal-students relationship.

Keywords: Females, Performance, Secondary school, Social anxiety Disorder, Students

INTRODUCTION

Social phobia is defined as a persistent and excessive fear of embarrassment in social situations when a person feels as if he/she is being observed by others and negatively evaluated [1]. According to the WHO, 27% of the adult population had a psychiatric disorder including depression, anxiety, and drug abuse that causes disability in their lives. Anxiety disorders rank sixth and account for 4% of all disabilities [2]. SAD is one of the most common types of anxiety disorders worldwide and was estimated to be 7% in the US [1, 3]. It is more common among females and among those who had low educational and socioeconomic status as well as among those who reported unhealthy parenting styles [4-6]. It was found that if not diagnosed and treated early, SAD can lead to other mental diseases like depression and alcohol and substance abuse [3, 7]. The prevalence of SAD in Saudi Arabia ranged from 11.7% to 25.8% and was associated significantly with parenting style [6, 8]. Other previous studies from Ethiopia, Kurdistan, and Turkey found that the prevalence of SAD was 27.5%, 31.25%, and 20.9% respectively and SAD was higher among females [9-11].

This study aimed to measure the prevalence of SAD and its impact on academic performance among female secondary school students in Medina, Saudi Arabia.

MATERIALS AND METHODS

Study Setting and Sample

This observational analytical cross-sectional study was conducted among 405 female secondary school students in Medina city, Saudi Arabia. Al Madinah was divided into four regions and two schools were selected randomly from each region. All students in the 8 governmental schools were approached.

Study Instruments

A self-administered questionnaire consisting of three parts was used in this study. The first part included questions on sociodemographic characteristics like age, school grade, family size, birth order, parental marital status, family income, and parenting style. The second part assessed SAD by using the Social Phobia Inventory (SPIN). The SPIN is a short, self-rating scale developed by Dr. K.M. Connor to capture social phobia symptoms [12-14]. It consists of 17 items and each item is rated from 0 (not at all) to 4 (extremely) with a total score of 0-68. A score ≥ 19 suggests SAD. It has good test-retest reliability, internal consistency, and convergent and divergent validity. The Cronbach alpha was 0.85 [15, 16]. The third part was the Liebowitz Social Anxiety Scale (LSAS) which is used to assess the severity of SAD. The LSAS is a self-rating scale developed by Dr. Michael Liebowitz to rate fear/anxiety and avoidance regarding 24 commonly feared performance or social situations [17]. It consists of 13 performance-related items and 11 social-related items which are rated from 0 (none/never) to 3 (severe/usually). The Cronbach alpha for the fear/anxiety domain was 0.88 and for the avoidance, the domain was 0.87 [15]. A score of <55 suggests mild SAD, 55-64 suggests moderate SAD, 65-79 suggests marked SAD, 80-94 suggests severe SAD, and >95 suggests very severe SAD. To evaluate the academic performance of the students, the last semester score of each student was obtained from the school.

Statistical Analysis

Analysis was performed using the Statistical Package for the Social Sciences (SPSS®) (version 22.0, IBM, Armonk, NY). Percentages and frequencies were obtained for the categorical variables. A Chi-square test was performed to assess the association between the categorical variables. The accepted level of significance was set below 0.05 ($p < 0.05$).

Ethical Considerations

Ethical approval was obtained from the Ethics Committee of the Directorate of Health in Al-Madinah. The objectives and benefits of the study were explained to the participants. Participants' confidentiality and anonymity were assured. Signed consents were obtained from the participants.

RESULTS

The study included 405 female high school students. The age of 41.8% of them was ≥ 18 years whereas 58.2% were aged ≤ 17 years. About half of them (48.1%) were in the third grade. The majority of them (84%) live with both parents whereas 16% live with one parent. The family size without counting parents ranged between 4 and 8 persons among 70.9% of the students. The first birth order was mentioned by 25.7% of them. Higher educated mothers and fathers (university) were reported among 32.6% and 38.5%, respectively. Family monthly income of less than 6000 SR was reported by 51.3% of the students. Regarding performance, almost half of them (50.1%) got $>90\%$ in the last semester (Table 1).

Table 1 Socio-demographic characteristics of the participants (n=405)

| Variables | | n | % |
|----------------------------|--------------|-----|-------|
| Age (Years) | ≤ 17 | 236 | 58.2% |
| | ≥ 18 | 169 | 41.8% |
| Educational grade | First | 87 | 21.5% |
| | Second | 123 | 30.4% |
| | Third | 195 | 48.1% |
| Living status | Both parents | 340 | 84% |
| | One parent | 65 | 16% |
| Family size | <4 | 54 | 13.3% |
| | 4-8 | 287 | 70.9% |
| | >8 | 64 | 15.8% |
| Birth order | First | 104 | 25.7% |
| | 2-3 | 141 | 34.8% |
| | 4-6 | 97 | 24% |
| | >6 | 63 | 15.5% |
| Maternal educational level | Illiterate | 16 | 4% |
| | Read/write | 29 | 7.2% |
| | School | 228 | 56.3% |
| | University | 132 | 32.6% |
| Paternal educational level | Illiterate | 10 | 2.5% |
| | Read/write | 21 | 5.2% |
| | School | 218 | 53.8% |

| | | | |
|----------------------------------|------------|-----|-------|
| | University | 156 | 38.5% |
| Family income (SAR/month) | ≤ 6000 | 208 | 51.3% |
| | 6001-10000 | 110 | 27.2% |
| | >10000 | 87 | 21.5% |
| Performance in the last semester | ≤ 80% | 75 | 18.5% |
| | 81%-90% | 127 | 31.4% |
| | >90% | 203 | 50.1% |

Relationship with Parents

Parental anger was usually reported by 27.2% of the students. Usual parental criticism was reported by 23% of them and it was in front of others in 8.4% of cases. Usual parental over-protection was reported by 59% of them. Parental provocation was usual among 16% of the students. Parental abuse was reported by 27.4% of the students. The prevalence of parental physical, and emotional abuse and parental neglect were 11.4%, 24.2%, and 8.6% respectively (Table 2).

Table 2 Students' relationship with their parents

| | N | % |
|--|-----|-------|
| Parental anger | | |
| Usually | 110 | 27.2% |
| Rarely | 211 | 52.1% |
| Never | 84 | 20.7% |
| Parental criticism | | |
| Usually | 93 | 23% |
| Rarely | 190 | 46.9% |
| Never | 122 | 30.1% |
| Parental criticism in front of others | | |
| Usually | 34 | 8.4% |
| Rarely | 112 | 27.7% |
| Never | 295 | 63.9% |
| Parental over-protection | | |
| Usually | 239 | 59.0% |
| Rarely | 110 | 27.2% |
| Never | 56 | 13.8% |
| Parental provocation | | |
| Usually | 65 | 16.0% |
| Rarely | 122 | 30.1% |
| Never | 218 | 53.8% |
| Parental abuse | | |
| No | 294 | 72.6% |
| Yes | 111 | 27.4% |
| Parental physical abuse | | |
| No | 359 | 88.6% |
| Yes | 46 | 11.4% |

| Parental emotional abuse | | |
|--------------------------|-----|-------|
| No | 307 | 75.8% |
| Yes | 98 | 24.2% |
| Parental neglect | | |
| No | 370 | 91.4% |
| Yes | 35 | 8.6% |

Prevalence and Severity of SAD

Table 3 summarizes the response of the participants to the individual items of the Social Phobia Inventory scale. Overall, the prevalence of social anxiety disorder among female high school students was 51.1%. Marked, severe, and very severe SAD were reported by 12.8%, 4.7%, and 3.7% respectively as obtained from the LSAS questionnaire.

Table 3 Response of the students to the Social Phobia Inventory scale

| | Not At All | A Little Bit | Somewhat | Very Much | Extremely |
|---|-------------|--------------|-------------|------------|------------|
| | N (%) | N (%) | N (%) | N (%) | N (%) |
| I am afraid of people in authority. | 169 (41.7%) | 99 (24.4%) | 86 (21.2%) | 37 (9.1%) | 14 (3.5%) |
| I am bothered by blushing in front of people. | 160 (39.5%) | 86 (21.2%) | 84 (20.7%) | 48 (11.9%) | 27 (6.7%) |
| Parties and social events scare me. | 216 (53.3%) | 81 (20.0%) | 52 (12.8%) | 39 (9.6%) | 17 (4.2%) |
| I avoid talking to people I don't know. | 92 (22.7%) | 90 (22.2%) | 108 (26.7%) | 78 (19.3%) | 37 (9.1%) |
| Being criticized scares me a lot. | 110 (27.2%) | 90 (22.2%) | 73 (18.0%) | 70 (17.3%) | 62 (15.3%) |
| I avoid doing things or speaking to people for fear of embarrassment. | 167 (41.2%) | 86 (21.2%) | 74 (18.3%) | 43 (10.6%) | 35 (8.6%) |
| Sweating in front of people causes me distress. | 217 (53.6%) | 78 (19.3%) | 48 (11.9%) | 33 (8.1%) | 29 (7.2%) |
| I avoid going to parties. | 218 (53.8%) | 67 (16.5%) | 59 (14.6%) | 37 (9.1%) | 24 (5.9%) |
| I avoid activities in which I am the centre of attention. | 197 (48.6%) | 85 (21.0%) | 56 (13.8%) | 40 (9.9%) | 27 (6.7%) |
| Talking to strangers scares me. | 194 (47.9%) | 88 (21.7%) | 61 (15.1%) | 35 (8.6%) | 27 (6.7%) |
| I avoid having to give speeches. | 123 (30.4%) | 72 (17.8%) | 72 (17.8%) | 67 (16.5%) | 71 (17.5%) |
| I would do anything to avoid being criticized. | 114 (28.1%) | 93 (23.0%) | 60 (14.8%) | 69 (17.5%) | 69 (17.0%) |
| Heart palpitations bother me when I am around people. | 177 (43.7%) | 79 (19.5%) | 55 (13.6%) | 50 (12.3%) | 44 (10.9%) |
| I am afraid of doing things when people might be watching. | 151 (37.3%) | 83 (20.5%) | 60 (14.8%) | 68 (16.8%) | 43 (10.6%) |
| Being embarrassed or looking stupid are among my worst fears. | 151 (37.3%) | 114 (28.1%) | 63 (15.6%) | 39 (9.6%) | 38 (9.4%) |
| I avoid speaking to anyone in authority. | 211 (52.1%) | 78 (19.3%) | 61 (15.1%) | 38 (9.4%) | 17 (4.2%) |
| Trembling or shaking in front of others is distressing to me. | 155 (38.3%) | 95 (23.5%) | 57 (14.1%) | 51 (12.6%) | 47 (11.6%) |

Association between Social Anxiety Disorder and Socio-Demographic Factors

SAD was higher among first-order students (60.6%), ($p=0.038$), among students whose fathers were illiterates (80%), ($p=0.041$) and among those who had low income (55.8%), ($p=0.022$) (Table 4).

Table 4 Association between socio-demographic factors and social anxiety disorder among participants

| | Social Anxiety Disorder | | p-value |
|------------------------------------|-------------------------|-------------|---------|
| | Yes N (%) | No N(%) | |
| Age (Years) | | | |
| ≤ 17 | 125 (53%) | 111 (47%) | 0.466 |
| ≥ 18 | 82 (48.5%) | 87 (51.5%) | |
| Educational grade | | | |
| First | 47 (54.0%) | 40 (46.0%) | 0.821 |
| Second | 56 (45.5%) | 67 (54.5%) | |
| Third | 104 (53.3%) | 91 (46.7%) | |
| Living status | | | |
| With parents | 168 (49.4) | 172 (50.6) | 0.083 |
| One parent | 39 (60) | 26 (40) | |
| Family size | | | |
| <4 | 26 (48.1%) | 28 (51.9%) | 0.697 |
| 4-8 | 152 (53.0%) | 135 (47.0%) | |
| >8 | 29 (45.3%) | 35 (54.7%) | |
| Birth order | | | |
| First | 63 (60.6%) | 41 (39.4%) | 0.038 |
| 2-3 | 61 (43.3) | 80 (56.7%) | |
| 4-6 | 47 (48.5%) | 50 (51.5%) | |
| >6 | 36 (57.1%) | 27 (42.9%) | |
| Maternal educational level | | | |
| Illiterate | 11 (68.8%) | 5 (31.2%) | 0.464 |
| Read/write | 15 (51.7%) | 14 (48.3%) | |
| School | 116 (50.9%) | 112 (49.1%) | |
| University | 65 (49.2%) | 67 (50.8%) | |
| Paternal educational level | | | |
| Illiterate | 8 (80.0%) | 2 (20.0%) | 0.041 |
| Read/write | 13 (61.9%) | 8 (38.1) | |
| School | 113(51.8%) | 105 (48.2%) | |
| University | 73 (46.8%) | 83 (53.2%) | |
| Family income (SR/month) | | | |
| ≤ 6000 | 116 (55.8%) | 92 (44.2%) | 0.022 |
| 6001-10000 | 51 (46.4%) | 59 (53.6%) | |
| >10000 | 40 (46.0%) | 47 (54.0%) | |
| Degree in the last semester | | | |
| ≤ 80% | 37 (49.3%) | 38 (50.7%) | 0.939 |
| 81%-90% | 65 (51.2%) | 62 (48.8%) | |
| >90% | 105 (51.7%) | 98 (48.3%) | |

Association between Social Anxiety Disorder and Parental-Students Relationship

SAD was higher among those who reported their parents were angry with them usually ($p<0.001$), those who reported their parents criticized them usually ($p<0.001$), and those who reported their parents criticized them in

front of others ($p < 0.001$) and those who reported usual parental provocation ($p = 0.005$). SAD was also higher among those who reported parental physical abuse ($p = 0.019$), emotional abuse ($p = 0.006$), and neglect ($p = 0.031$) (Table 5).

Table 5 Association between social anxiety disorder and parental-students relationship among participants

| | Social Anxiety Disorder | | p-value |
|--|-------------------------|-------------|---------|
| | Yes N (%) | No N (%) | |
| | N=207 | N=198 | |
| Parental anger | | | |
| Usually | 69 (62.7%) | 41 (37.3%) | <0.001 |
| Rarely | 109 (51.7%) | 102 (48.3%) | |
| Never | 29 (34.5%) | 55 (65.5%) | |
| Parental criticism | | | |
| Usually | 64 (68.8%) | 29 (31.2%) | <0.001 |
| Rarely | 94 (49.5%) | 96 (50.5%) | |
| Never | 49 (40.2%) | 73 (59.8%) | |
| Parental criticism in front of others | | | |
| Usually | 21 (61.8%) | 13 (38.2%) | <0.001 |
| Rarely | 74 (66.1%) | 38 (33.9%) | |
| Never | 112 (43.2%) | 147 (56.8%) | |
| Parental over-protection | | | |
| Usually | 123 (51.5%) | 116 (48.5%) | 0.98 |
| Rarely | 56 (50.9%) | 54 (49.1%) | |
| Never | 28 (50.0%) | 28 (50.0%) | |
| Parental provocation | | | |
| Usually | 42 (64.6%) | 23 (35.4%) | 0.005 |
| Rarely | 69 (56.6%) | 53 (43.4%) | |
| Never | 96 (44.0%) | 122 (56.0%) | |
| Parental abuse | | | |
| No | 134 (45.6%) | 160 (54.4%) | <0.001 |
| Yes | 73 (65.8%) | 38 (34.2%) | |
| Parental physical abuse | | | |
| No | 176 (49.0%) | 183 (51.0%) | 0.019 |
| Yes | 31 (67.4%) | 15 (32.6%) | |
| Parental emotional abuse | | | |
| No | 145 (47.2%) | 162 (52.8%) | 0.006 |
| Yes | 62 (63.3%) | 36 (36.7%) | |
| Parental neglect | | | |
| No | 183 (49.5%) | 187 (50.5%) | 0.031 |
| Yes | 24 (68.6%) | 11 (31.4%) | |

DISCUSSION

The present study aimed to assess the prevalence and severity of SAD and determine its associated factors as well as its impact on the academic performance of secondary school girls in Medina city, Saudi Arabia [18,19]. In the present study, the prevalence of SAD among secondary school girls was 51.1%. Regarding its severity, it was mild among 31.4% and moderate among 17% of the students. Marked, severe and very severe form was reported by 21.2% of the students. The prevalence of SAD ranged from 6.2% to 32.4% in some Middle East countries, 14.1% to 10.3% in Asian countries, and 26.5% in Poland [20-30]. Moreover, in an earlier comparative study, the prevalence of social phobia in Saudi Arabia, the United Arab of Emirates, and Egypt were 9.8%, 7.8%, and 13%, respectively [30]. In a recent study carried out among undergraduate students at Jazan University, Saudi Arabia, the prevalence of SAD was 25.8%; it was mild in 47.2% of them, moderate, marked, severe/very severe in 42.3% and 10.5% of them, respectively [7]. The apparent difference in the rate of SAD between different studies could be related to some reasons; the demographic characteristics of the participants particularly age, gender, socio-cultural variations between different countries, and the utilization of different tools to assess social phobia.

Relatively, few studies investigated social phobia in only one sex as a result of cultural issues and all were done in Saudi Arabia, as in the present study. For example, the prevalence of SAD among secondary school boys was 11.7% in Abha city and 14.1% in Khamis Mushait [8, 23]. In Taif, the prevalence rate among female university students was 16.3% [31].

In agreement with others, the present study revealed that SAD was more reported among first-birth order students than others [32-34], However, others reported that social phobia was more among children born later [35].

Lower-educated fathers and lower family income, which indicate low socioeconomic status, were associated with a higher rate and more severe form of SAD among female students in the present study. This is quite expected as it is known that low socioeconomic status is associated with many psychological problems including anxiety disorders, as low socioeconomic status induces psychological problems, which in turn may cause individuals to become of low socioeconomic status [36,37].

Additionally, higher-grade students were more likely to have severe/very severe forms of SAD than their counterparts. However, Al-Qahtani didn't find a relation between school grades and the prevalence and severity of social phobia [32].

Regarding the parental-student relationship, usual parental anger toward students, criticizing them individually or in front of others, parental provocation, and parental abuse of all types (physical, emotional, and neglect) were associated with a higher rate of SAD. In addition, a more severe form of SAD was associated with usual parental criticism of students either individually or in front of others, parental provocation, and parental abuse particularly, physical type. Similar findings were reported in two previous studies [8, 38]. Bracik J, *et al.* found that the environmental factors of the school and family had an impact on the development of SAD in adolescents [23].

The present study and others didn't observe an association between the SAD and self-reported academic performance of the students [10, 39]. In the present study, academic performance was self-rated, Therefore further investigation using academic performance from students' records is recommended. Two main limitations are present in this study. The first limitation is the inclusion of females only, which can't give an overview of the problem among the adolescent population in Medina. The second limitation is the cross-

sectional design of the study which can't show the causal relationship between SAD and possible related factors. Despite those limitations, this study could be of usefulness to decision-makers as this is a hidden problem in our conservative society, particularly among female adolescents.

CONCLUSION

SAD is a common health problem among girls enrolled in secondary schools in Medina city as about half of them reported the problem. Although it was mainly mild in severity, a considerable proportion had severe/very severe levels. It was more reported among first birth order students and those with lower socioeconomic status. The parental style was associated with a higher rate of SAD.

Recommendations

Initial and regular periodic evaluations of secondary school girls in Medina city for SAD should be done to discover early enough cases. Health education is recommended for both parents and teachers regarding possible risk factors and early symptoms of social phobia to increase their awareness of the bad impacts of physical, and emotional abuse and their effects on students' psychological health. Special attention should be given to students of low socioeconomic status, by providing more social and psychological support. Future studies are recommended to include male students and those in earlier school stages.

DECLARATIONS

Conflicts of Interest

The authors declared no potential conflicts of interest concerning the research, authorship, and/or publication of this article.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

REFERENCES

- [1] American Psychiatric Association, A. "Diagnostic and statistical manual of mental disorders." Vol. 3, Washington, DC: American Psychiatric Association, 1980.
- [2] Murray, Christopher JL, et al. "The Global Burden of Disease 2000 project: Aims, methods and data sources." *Geneva: World Health Organization*, Vol. 36, 2001, pp. 1-57.
- [3] Mekuria, Kindie, et al. "High magnitude of social anxiety disorder in school adolescents." *Psychiatry Journal*, Vol. 2017, 2017.
- [4] Spence, Susan H., and Ronald M. Rapee. "The aetiology of social anxiety disorder: An evidence-based model." *Behaviour Research and Therapy*, Vol. 86, 2016, pp. 50-67.
- [5] Fors, Payton Q., and Deanna M. Barch. "Differential relationships of child anxiety and depression to child report and parent report of electronic media use." *Child Psychiatry & Human Development*, Vol. 50, No. 6, 2019, pp. 907-17.
- [6] Ghazwani, Jaafar Y., Shamsun N. Khalil, and Razia A. Ahmed. "Social anxiety disorder in Saudi adolescent boys: Prevalence, subtypes, and parenting style as a risk factor." *Journal of Family & Community Medicine*, Vol. 23, No. 1, 2016, pp. 25-31.

- [7] Stein, Murray B., and Dan J. Stein. "Social anxiety disorder." *The Lancet*, Vol. 371, No. 9618, 2008, pp. 1115-25.
- [8] Hakami, Ramzi M., et al. "Social anxiety disorder and its impact in undergraduate students at Jazan University, Saudi Arabia." *Mental Illness*, Vol. 9, No. 2, 2017, pp. 42-47.
- [9] Stein, Dan J., et al. "The cross-national epidemiology of social anxiety disorder: Data from the World Mental Health Survey Initiative." *BMC Medicine*, Vol. 15, No. 1, 2017, pp. 1-21.
- [10] Gardi, Abdulqader H.H. "Prevalence of social phobia among high school students in Erbil, Kurdistan region." *Zanco Journal of Medical Sciences*, Vol. 20, No. 3, 2016, pp. 1497-504.
- [11] Gultekin, Bulent K., and Ferhan I. Dereboy. "The prevalence of social phobia, and its impact on quality of life, academic achievement, and identity formation in university students." *Turkish Journal of Psychiatry*, Vol. 22, No. 3, 2011, pp. 150-58.
- [12] Education M. Saudi Arabia-Ministry of Education, 2018. <https://www.moe.gov.sa/ar/Pages/StatisticalInformation.aspx>
- [13] Raosoft. "Sample size calculator." 2019. <http://www.raosoft.com/samplesize.html>
- [14] Connor, Kathryn M., et al. "Psychometric properties of the Social Phobia Inventory (SPIN): New self-rating scale." *The British Journal of Psychiatry*, Vol. 176, No. 4, 2000, pp. 379-86.
- [15] Shah, P., and Lakhan Kataria. "Social phobia and its impact in Indian university students." *The Internet Journal of Mental Health*, Vol. 6, No. 2, 2010, pp. 1-8.
- [16] Poormohammad, Ahmad, et al. "Social phobia in persian adults with stuttering." *Journal of Rehabilitation Sciences & Research*, Vol. 5, No. 1, 2018, pp. 13-17.
- [17] Liebowitz, Michael R. "Social phobia." *Modern Problems of Pharmacopsychiatry*, Vol. 22, 1987, pp. 141-73.
- [18] Charan, Jaykaran, and Tamoghna Biswas. "How to calculate sample size for different study designs in medical research?" *Indian Journal of Psychological Medicine*, Vol. 35, No. 2, 2013, pp. 121-26.
- [19] Crome, Erica, Andrew Baillie, and Alan Taylor. "Are male and female responses to social phobia diagnostic criteria comparable?" *International Journal of Methods in Psychiatric Research*, Vol. 21, No. 3, 2012, pp. 222-31.
- [20] Hummadi, Basheer F., and Abdulkareem Alobaidi. "Social phobia among secondary school students in Baghdad/Iraq." *Journal of the Canadian Academy of Child and Adolescent Psychiatry*, Vol. 23, No. 1, 2014, pp. 70-71.
- [21] Hussein, Ahmed H., et al. "Social phobia among secondary school students in Babil, Iraq." *Journal of the Faculty of Medicine Baghdad*, Vol. 58, No. 4, 2016, pp. 354-56.
- [22] Jefferies, Philip, and Michael Ungar. "Social anxiety in young people: A prevalence study in seven countries." *PLoS One*, Vol. 15, No. 9, 2020, p. e0239133.
- [23] Bracik, Joanna, Krzysztof Krysta, and Adam Zaczek. "Impact of family and school environment on the development of social anxiety disorder: A questionnaire study." *Danubina Psychiatry*, Vol. 24, No. 1, 2012, pp. 125-27.
- [24] Mazhari, Shahrzad, Maryam Ekhlaspour, and Nabi Banazadeh. "Social phobia and its association with academic performance among student of Kerman University of Medical Sciences Iran." *Strides in Development of Medical Education*, Vol. 11, No. 2, 2014, pp. 227-35.

- [25] Momani, F., and A. M. Jaradat. "Social phobia among university students: Prevalence and socio-demographic factors." *Jordan Journal of Social Sciences*, Vol. 4, No. 1, 2011, pp. 71-88.
- [26] Jin, Yuelong, et al. "Prevalence and risk factors of anxiety status among students aged 13-26 years." *International Journal of Clinical and Experimental Medicine*, Vol. 7, No. 11, 2014, pp. 4420-26.
- [27] Srinivasan, Manikandan, et al. "Comment on: Social anxiety disorder in Saudi adolescent boys: Prevalence, subtypes, and parenting styles as a risk factor." *Journal of Family & Community Medicine*, Vol. 23, No. 3, 2016, pp. 185-86.
- [28] Burstein, Marcy, et al. "Social phobia and subtypes in the National Comorbidity Survey-Adolescent Supplement: Prevalence, correlates, and comorbidity." *Journal of the American Academy of Child & Adolescent Psychiatry*, Vol. 50, No. 9, 2011, pp. 870-80.
- [29] Dodangi, Nasrin, Nastaran Habibi Ashtiani, and Burhanoddin Valadbeigi. "Prevalence of DSM-IV TR psychiatric disorders in children and adolescents of Paveh, a western city of Iran." *Iranian Red Crescent Medical Journal*, Vol. 16, No. 7, 2014, p. e16743.
- [30] Azab, Hemaïd M. Musstafa, Usama A. El-Khouli, and A. Magdi. "Social phobia among secondary school students: A comparative study between Egypt, Saudi Arabia and the United Arab Emirates." *Egyptian Journal of Psychiatry*, Vol. 26, No. 1, 2007.
- [31] Taha, Azza Ali, et al. "Social anxiety disorder and its correlates among female students at Taif University, Saudi Arabia." *Research in Psychology and Behavioral Sciences*, Vol. 5, No. 2, 2017, pp. 50-56.
- [32] Al-Qahtani, A. M., and A. B. F. M. Sbfm. "Prevalence and risk factors of social phobia among secondary school male students in Khamis Mushayt, Kingdom of Saudi Arabia." *The Medical Journal of Cairo University*, Vol. 80, No. 1, 2012, pp. 871-76.
- [33] Voci, Sabrina C., et al. "Social anxiety in late adolescence: The importance of early childhood language impairment." *Journal of Anxiety Disorders*, Vol. 20, No. 7, 2006, pp. 915-30.
- [34] Aljohani, Basem Musllam, and Amani Mahrus. "Prevalence and possible risk factor of social phobia among male secondary and intermediate school students in Al-Medinah, 2016." *The Egyptian Journal of Hospital Medicine*, Vol. 72, No. 7, 2018, pp. 4836-42.
- [35] Hudson, Jennifer L., and Ronald M. Rapee. "The origins of social phobia." *Behavior Modification*, Vol. 24, No. 1, 2000, pp. 102-29.
- [36] Bromet, Evelyn, et al. "Cross-national epidemiology of DSM-IV major depressive episode." *BMC Medicine*, Vol. 9, No. 1, 2011, pp. 1-16.
- [37] Kawakami, Norito, et al. "Early-life mental disorders and adult household income in the World Mental Health Surveys." *Biological Psychiatry*, Vol. 72, No. 3, 2012, pp. 228-37.
- [38] Abdallah, Eman Shokry, et al. "Association between social phobia and parenting styles among secondary school students." *American Journal of Nursing Science*, Vol. 5, No. 3, 2016, pp. 96-105.
- [39] Strahan, Esther Y. "The effects of social anxiety and social skills on academic performance." *Personality and Individual Differences*, Vol. 34, No. 2, 2003, pp. 347-66.